arXiv:math/0506275v2 [math.DG] 28 Jun 2005

THE GEOMETRIC THEORY OF THE FUNDAMENTAL GERM

T.M. GENDRON

ABSTRACT. The fundamental germ is a generalization of π_1, first defined for laminations which arise through group actions [4]. In this paper, the fundamental germ is extended to any lamination having a dense leaf admitting a smooth structure. In addition, an amplification of the fundamental germ called the mother germ is constructed, which is, unlike the fundamental germ, a topological invariant. The fundamental germs of the antenna lamination and the $PSL(2,\mathbb{Z})$ lamination are calculated, laminations for which the definition in [4] was not available. The mother germ is used to give a proof of a Nielsen theorem for the algebraic universal cover of a closed surface of hyperbolic type.

1. INTRODUCTION

This paper represents a continuation of our quest to extend $\mathbb{Z}$-coefficient algebraic topology to laminations through the generalization of π_1 called the fundamental germ. In this paper, we extend this construction to any lamination admitting a smooth structure.

Let us recall briefly the intuition behind the fundamental germ. Consider a suspension

$$\mathscr{L}_\rho \;=\; \left(\widetilde{B}\times \mathsf{T}\right)\Big/\pi_1 B$$

of a representation $\rho : \pi_1 B \to \mathsf{Homeo}(\mathsf{T})$, where B is a manifold. Then $\pi_1 B$ acts on $\mathscr{L}_\rho$ as fiber preserving homeomorphisms. Let $T \approx \mathsf{T}$ be a fiber transversal and let $x_0, x, \in T$. A $\pi_1 B$-diophantine approximation of $x \in T$ based at x_0 is a sequence $\{g_\alpha\} \subset \pi_1 B$ with $g_\alpha \cdot x_0 \to x$. The fundamental germ $[\![\pi]\!]_1(\mathscr{L}, x_0, x)$ is then the groupoid of tail equivalence classes of sequences of the form $\{g_\alpha \cdot h_\alpha^{-1}\}$ where $\{g_\alpha\}$, $\{h_\alpha\}$ are diophantine approximations of x along x_0 [4]. This construction is more generally available for any lamination occurring as a quotient of a suspension, a double-coset of a Lie group or a locally-free action of a Lie group on a space, laminations which we refer to collectively as *algebraic*. Intuitively, if L is the leaf containing x_0, the elements of $[\![\pi]\!]_1(\mathscr{L}, x_0, x)$ can be thought of as sequences of paths in L whose endpoints converge transversally to x. Such a sequence can be thought of as an ideal loop based at x that records an “asymptotic identification” within the leaf L.

For a linear foliation $\mathscr{F}_r$ of a torus by lines of slope r, the diophantine analogy is literal and $[\![\pi]\!]_1(\mathscr{F}, x_0, x)$ is the group of classical diophantine approximations of r. A manifold B is a supension of the trivial representation *i.e.* a lamination with a single leaf and fiber transversals that are points, which forces $x_0 = x$. Then all sequences in $\pi_1 B$ converge, and we find that $[\![\pi]\!]_1(B, x) = {}^*\pi_1(B, x) =$ the nonstandard fundamental group of B.

We now turn to the contents of this article. The algebraic definition of the fundamental germ just described, while amenable to calculation, has the following serious drawbacks:

Date: 25 May 2005.

1991 *Mathematics Subject Classification.* Primary, 57R30, 14B0.

Key words and phrases. fundamental germ, laminations, foliations, diophantine approximation, mapping class group.

(1) It is available only for the select family of algebraic laminations.
(2) It is an invariant only with respect to the special class of *trained* lamination homeomorphisms (*c.f.* [4]).

Addressing these flaws is the central theme of the present study. In the summary that follows, we shall assume for simplicity that all leaves are simply connected.

We begin with item (1). Let $\mathscr{L}$ be an arbitrary lamination admitting a smooth structure, let x_0, x be as above and denote by L the leaf containing x_0. Equip $\mathscr{L}$ with a leaf-wise riemannian metric that has continuous transverse variation. In this paper, we shall refer to such a lamination as *riemannian*. The new idea here is to use the leaf-wise geometry to represent – as sequences of isometries – the diophantine approximations which would make up $\llbracket\pi\rrbracket_1$. If L has constant curvature geometry, this prescription may be followed word-for-word. Fixing a transversal T containing x_0, x and a continuous section of orthonormal frames $\mathsf{f} = \{\mathsf{f}_y\}$, $y \in T$, we define a diophantine approximation of x to be a sequence $\{A_\alpha\}$ of isometries of L for which $(A_\alpha)_* \mathsf{f}_{x_0}$ belongs to f and converges transversally to f_x. The fundamental germ $\llbracket\pi\rrbracket_1(\mathscr{L}, x_0, x, \mathsf{f})$ is then defined to be the set of tails of sequences of the form $\left\{A_\alpha B_\alpha^{-1}\right\}$ where $\{A_\alpha\}$, $\{B_\alpha\}$ are diophantine approximations of x.

In the case of non constant curvature leaf-wise geometry, it is necessary to work within the category of *virtual geometry* in order to make sense of the notion of diophantine approximation. There, a riemannian manifold M is replaced by a union of riemannian manifolds, its virtual extension ${}^{\bullet}M$, which consists of all sequences in M up to the relation of being asymptotic. A virtual isometry between riemannian manifolds M and N consists of a pair of isometric inclusions ${}^{\bullet}M \leftrightarrows {}^{\bullet}N$. All dense leaves of a riemannian lamination have virtually isometric universal covers, and moreover, a dense leaf having no ordinary isometries will admit many virtual isometries.

This leads to the following definition of a diophantine approximation: let $x \in T$, f a frame field on T and let L be any leaf accumulating on x. Then a sequence $\mathsf{f}_{x_\alpha} \rightarrow \mathsf{f}_x$, $\{x_\alpha\} \subset L$, determines an isometry ${}^{\bullet}f : L_x \rightarrow U \subset {}^{\bullet}L$, where L_x is the leaf containing x and U is a component of ${}^{\bullet}L$. The fundamental germ $\llbracket\pi\rrbracket_1(\mathscr{L}, L, x, \mathsf{f})$ is defined to be the set of (maximal extensions of) maps of the form

$$ {}^{\bullet}f \circ {}^{\bullet}g^{-1}. $$

In this way, we now have a definition of the fundamental germ valid for any lamination admitting a smooth structure along the leaves.

In order to address drawback (2), we will need the *germ universal cover*

$$ \llbracket\widetilde{\mathscr{L}}\rrbracket \ \subset \ {}^{\bullet}L, $$

defined to be the set of asymptotic classes of sequences in L that converge to points of $\mathscr{L}$. The germ universal cover plays the role of a unit space for a groupoid structure on $\llbracket\pi\rrbracket_1(\mathscr{L}, L, x, \mathsf{f})$. It is a lamination whose leaves are nowhere dense, and when L is dense, the canonical map $\llbracket\widetilde{\mathscr{L}}\rrbracket \rightarrow \mathscr{L}$ is onto. We may therefore think of $\llbracket\widetilde{\mathscr{L}}\rrbracket$ as obtained from $\mathscr{L}$ by "unwrapping" all transversal topology implemented by L.

Assume now that L is dense. The *mother germ* $\llbracket\pi\rrbracket_1\mathscr{L}$ is defined to be the groupoid of all partially defined maps of $\llbracket\widetilde{\mathscr{L}}\rrbracket$ that are homeomorphisms on domains which are sublaminations of $\llbracket\widetilde{\mathscr{L}}\rrbracket$ and preserve the projection $\llbracket\widetilde{\mathscr{L}}\rrbracket \rightarrow \mathscr{L}$. We have in particular that

$$ \llbracket\pi\rrbracket_1\mathscr{L} \backslash \llbracket\widetilde{\mathscr{L}}\rrbracket \ \cong \ \mathscr{L}. $$

$\llbracket\pi\rrbracket_1\mathscr{L}$ is the receptacle of all the $\llbracket\pi\rrbracket_1(\mathscr{L}, L', x, \mathsf{f})$ for L' dense, in that it contains subgroupoids isomorphic to each. The mother germ is functorial with respect to topological lamination covering maps, and is therefore, in spite of its riemannian construction, a *topological* invariant. This takes care of item (2) above.

The remainder of the paper is devoted to examples and an application. Many examples were discussed in [4], and so for this reason we limit ourselves to laminations which are not algebraic and hence which do not have a fundamental germ in the sense described there.

The first example we consider is that which we call here the *antenna lamination*, a surface lamination discovered by Kenyon and Ghys [6] which has the distinction of having leaves of both parabolic and hyperbolic type. With respect to a hyperbolic leaf, the fundamental germ is calculated as a set to be ${}^*F_2 \times ({}^*\mathbb{Z}_{\hat{2}} \oplus {}^*\mathbb{Z}_{\hat{2}})$ where ${}^*F_2$ is the nonstandard free group on two generators, and ${}^*\mathbb{Z}_{\hat{2}}$ is the subgroup of ${}^*\mathbb{Z}$ isomorphic to the fundamental germ of the dyadic solenoid. Although a product of groups, this germ is not a group with respect to its defined multiplication. It is the first example we have encountered of a fundamental germ that is not a group.

The second example is that of the Anosov foliation of the unit tangent bundle to the modular surface. Although this is just the suspension of the action of $PSL(2,\mathbb{Z})$ on the boundary of the hyperbolic plane, the definition of the fundamental germ found in [4] is unavailable since it does not work for actions with fixed points. We calculate the fundamental germ here as a set to be $PSL(2,{}^*\mathbb{Z})$, but as in the case of the antenna lamination, it is also not a group with respect to its defined multiplication.

The final result of this paper concerns the use of the fundamental germ to calculate the mapping class group of the algebraic universal cover $\widehat{\Sigma}$ of a closed surface Σ of hyperbolic type. $\widehat{\Sigma}$ is by definition the inverse limit of finite covers of Σ, a compact solenoid with dense disk leaves. If $L \subset \widehat{\Sigma}$ is a fixed leaf, the leafed mapping class group $\mathsf{MCG}(\mathscr{L},L)$ is the quotient $\mathsf{Homeo}_+(\mathscr{L},L)/\simeq$, where $\mathsf{Homeo}_+(\mathscr{L},L)$ denotes the group of orientation-preserving homeomorphisms fixing set-wise L and $\simeq$ denotes homotopy. If we denote by $\mathsf{Vaut}(\pi_1\Sigma)$ the group of virtual automorphisms of $\pi_1\Sigma$ (*c.f.* §10) then

Theorem. *There is an isomorphism*

$$\Theta : \mathsf{MCG}(\mathscr{L},L) \longrightarrow \mathsf{Vaut}(\pi_1\Sigma).$$

A proof of this theorem first appeared in the unpublished 1997 thesis of C. Odden [8]. Due to its importance in the genus-independent expression of the Ehrenpreis conjecture [5], we provide a proof in order to ensure its inclusion in the literature.

Acknowledgements: I would like to thank P. Makienko and A. Verjovsky with whom I enjoyed fruitful conversations regarding several important aspects of this paper. I would also like to thank the Instituto de Matemáticas of the UNAM for providing a pleasant work enviroment and generous financial support.

2. Virtual Geometry

Virtual geometry is obtained as a quotient of nonstandard geometry, which we now review: references [7], [9], [10].

Let M be a topological space, $\mathfrak{U} \subset 2^{\mathbb{N}}$ an ultrafilter on the natural numbers all of whose elements have infinite cardinality. The *nonstandard space* ${}^*M$ is the set of sequences in M modulo $\mathfrak{U}$: that is,

$$\{x_i\} \sim \{y_i\} \text{ if and only if } \{x_i\}|_X = \{y_i\}|_X \text{ for some } X \in \mathfrak{U}. \tag{1}$$

Elements of ${}^*M$ are denoted ${}^*x$. There is a natural map $M \hookrightarrow {}^*M$ given by the constant sequences. Modulo the continuum hypothesis, ${}^*M$ is independent of the choice of ultrafilter.

There are two topologies on ${}^*M$ that naturally suggest themselves. The *enlargement topology* is generated by sets of the form ${}^*O$, where O is open in M. It has the same countability as the topology of M but is non-Hausdorff. The *internal topology* is generated by sets of the form $[O_\alpha] = \{{}^*x \in {}^*M \,|\, {}^*x \text{ is represented by a sequence } \{x_\alpha\},\ x_\alpha \in O_\alpha\}$, where $\{O_\alpha\}$ is any sequence of open sets of M. It is Hausdorff but has greater countability than the topology of M.

For example, if we let $M = \mathbb{R}$ we obtain the *nonstandard reals* ${}^*\mathbb{R}$, a totally ordered, non-archemidean field. Note that ${}^*\mathbb{R}$ is an infinite-dimensional vector space over $\mathbb{R}$. We will refer to the following substructures of the nonstandard reals:

- The subring of bounded nonstandard reals, denoted ${}^*\mathbb{R}_{\mathsf{fin}}$, which consists of all classes of sequences that are bounded.
- The additive subgroup of infinitesimals, denoted ${}^*\mathbb{R}_\varepsilon$, which consists of all classes of sequences converging to 0.
- The cone of positive elements, denoted ${}^*\mathbb{R}_+$, which consists of all classes of sequences that are ≥ 0.

${}^*\mathbb{R}_{\mathsf{fin}}$ is a local topological ring in either the enlargement or internal topology, with maximal ideal ${}^*\mathbb{R}_\varepsilon$. The quotient ${}^*\mathbb{R}_{\mathsf{fin}}/{}^*\mathbb{R}_\varepsilon$ is isomorphic to $\mathbb{R}$, homeomorphic with the quotient enlargement topology (the quotient internal topology is discrete). The inclusion $\mathbb{R} \hookrightarrow {}^*\mathbb{R}_{\mathsf{fin}}$ allows us to canonically identify ${}^*\mathbb{R}_{\mathsf{fin}}$ with the product $\mathbb{R} \times {}^*\mathbb{R}_\varepsilon$. Taking the product of the euclidean topology on $\mathbb{R}$ with the discrete topology on ${}^*\mathbb{R}_\varepsilon$, we obtain a third topology on ${}^*\mathbb{R}_{\mathsf{fin}}$ which is Hausdorff and quotients by ${}^*\mathbb{R}_\varepsilon$ to the topology on $\mathbb{R}$. We call this third topology the *lamination topology*: it may be extended to ${}^*\mathbb{R}$ by giving the group ${}^*\mathbb{R}/\mathbb{R}$ the discrete topology and identifying ${}^*\mathbb{R} \cong \mathbb{R} \times ({}^*\mathbb{R}/\mathbb{R})$.

If M is an n-manifold, then ${}^*M$ is a nonstandard manifold modelled on ${}^*\mathbb{R}^n$. If we denote by ${}^*M_{\mathsf{fin}}$ the points of ${}^*M$ represented by sequences which converge to points of M, then we may choose an atlas on ${}^*M_{\mathsf{fin}}$ whose transitions preserve the lamination structure of ${}^*\mathbb{R}^n_{\mathsf{fin}}$ *i.e.* ${}^*M_{\mathsf{fin}}$ is an n-lamination. In general, ${}^*M$ is a union of laminations of dimensions $\leq n$, this because of the possibility of "dimension collapse" which we describe in the proof of Theorem 1 below.

If d is a metric inducing the topology of M, it extends to a ${}^*\mathbb{R}_+$-valued metric ${}^*d$ on ${}^*M$. Write ${}^*x \simeq {}^*x'$ if ${}^*d({}^*x, {}^*x') \in {}^*\mathbb{R}_\varepsilon$.

Definition 1. The **virtual extension** of M is the quotient

$$ {}^\bullet M \;=\; {}^*M/\simeq, $$

equipped with the quotient lamination topology.

The virtual extension of ${}^\bullet\mathbb{R}$ of $\mathbb{R}$ is called the *virtual reals*, a totally-ordered real vector space. The metric ${}^*d$ on ${}^*M$ induces a ${}^\bullet\mathbb{R}_+$-valued metric ${}^\bullet d$ on ${}^\bullet M$. Given ${}^\bullet x \in {}^\bullet M$, the set

$$ U_{{}^\bullet x} \;=\; \{{}^\bullet y \,|\, {}^\bullet d({}^\bullet x, {}^\bullet y) \in \mathbb{R}\} $$

is a component of ${}^\bullet M$ called the *galaxy* of ${}^\bullet x$. M is a galaxy of ${}^\bullet M$, and ${}^\bullet M$ is the union of all of its galaxies.

The galaxies of ${}^\bullet M$ can be quite different from one another. For example if M is simply connected, then there may be galaxies that are not. For example, suppose that M is a noncompact leaf of the Reeb foliation of the torus. Consider a sequence of points $\{x_\alpha\}$ in M converging to a point $\hat{x}$ in the compact toral leaf. Let $\{\gamma_\alpha\}$ be a sequence of simple closed curves converging to the meridian through $\hat{x}$. Then the limit curve ${}^\bullet\gamma$ is essential in the universe $U_{{}^\bullet x}$. On the other hand, if M is a riemannian homogeneous space, then the universes of ${}^\bullet M$ are all isometric to M.

Theorem 1. *If M is a complete riemannian manifold of dimension n, each galaxy U of ${}^\bullet M$ has the structure of a complete riemannian manifold of dimension $m \leq n$.*

Proof. Given a galaxy U, ${}^\bullet x \in U$ and $\{x_\alpha\}$ a representative sequence, let m be the largest integer for which there exists a sequence of m-dimensional balls $\{D_r(x_\alpha)\}$ of fixed radius r about $\{x_\alpha\}$. The integer m is independent of the representative sequence and defines an m-ball $D_r({}^\bullet x) \subset U$. The function ${}^\bullet x \mapsto m$ is locally constant, thus the collection of such balls defines on U the structure of a smooth m-manifold. Note that it is possible to have $m < n$: for example, if M is a hyperbolic manifold with a cusp, then for a class of sequence emptying into the cusp, we have $m = n-1$.

Consider the nonstandard tangent bundle

$$\mathbf{T}^*M \ := \ {}^*\big(\mathbf{T}M\big).$$

There is a natural projection of $\mathbf{T}^*M$ onto ${}^*M$ whose fiber $\mathbf{T}_{{}^*x}{}^*M$ – the tangent space at ${}^*x$ – consists of classes of sequences of vectors $\{\mathsf{v}_\alpha\}$ based at sequences $\{x_\alpha\}$ belonging to the class of ${}^*x$. It is not difficult to see that $\mathbf{T}_{{}^*x}{}^*M$ is a real infinite-dimensional vector space. The riemannian metric ρ extends to a ${}^*\mathbb{R}$-valued metric ${}^*\rho$ on $\mathbf{T}^*M$ in the obvious way. Denote by ${}^*|\cdot|$ the associated norm. Define the bounded tangent bundle by

$$\mathbf{T}_{\mathsf{fin}}{}^*M \ = \ \Big\{ {}^*\mathsf{v} \in \mathbf{T}^*M \ \Big| \ {}^*|{}^*\mathsf{v}| \in {}^*\mathbb{R}_{\mathsf{fin}} \Big\}.$$

Given tangent vectors ${}^*\mathsf{v}$ and ${}^*\mathsf{v}'$ based at ${}^*x$ and ${}^*x'$, we write ${}^*\mathsf{v} \simeq {}^*\mathsf{v}'$ if

1. ${}^*x \simeq {}^*x'$.
2. the Levi-Civita parallel translate of a representative $\{\mathsf{v}_\alpha\}$ of ${}^*\mathsf{v}$ to a representative $\{x'_\alpha\}$ of ${}^*x'$ – along a sequence of geodesics connecting to a representative $\{x_\alpha\}$ of ${}^*x$ – is asymptotic to a representative $\{\mathsf{v}'_\alpha\}$ of ${}^*\mathsf{v}'$.

Now define the bounded tangent bundle of ${}^\bullet M$ to be

$$\mathbf{T}_{\mathsf{fin}}{}^\bullet M \ = \ \mathbf{T}_{\mathsf{fin}}{}^*M\big/\simeq .$$

The nonstandard riemannian metric ${}^*\rho$ on $\mathbf{T}_{\mathsf{fin}}{}^*M$ descends to a riemannian metric on $\mathbf{T}_{\mathsf{fin}}{}^\bullet M$. If U is a galaxy, its tangent space may be identified with the restriction of $\mathbf{T}_{\mathsf{fin}}{}^\bullet M$ to U. Now any geodesic $\eta \subset U$ can be realized as a sequence class of geodesics $\{\eta_\alpha\}$. Since each member of such a sequence can be continued indefinitely, the same is true of η, hence U is complete. $\square$

Definition 2. Let M, N be riemannian n-manifolds. A **virtual subisometry** is an injective map

$${}^\bullet f : {}^\bullet M \hookrightarrow {}^\bullet N,$$

where ${}^\bullet f$ maps each galaxy of ${}^\bullet M$ isometrically onto a galaxy of ${}^\bullet N$. If in addition there exists a virtual subisometry ${}^\bullet g : {}^\bullet N \hookrightarrow {}^\bullet M$, then the pair $({}^\bullet f, {}^\bullet g)$ is called a **virtual isometry**.

We write $M \leq_{\mathsf{vir}} N$ to indicate the existance of a virtual subisometry ${}^\bullet f$ and $M \cong_{\mathsf{vir}} N$ indicates the existence of a virtual isometry. The relation $\leq_{\mathsf{vir}}$ defines a partial ordering on the set of all riemannian n-manifolds.

An isometry $f : M \to N$ clearly induces a virtual isometry $({}^\bullet f, {}^\bullet g) : {}^\bullet M \leftrightarrows {}^\bullet N$ with ${}^\bullet f, {}^\bullet g$ inverse to one another. More generally, a continuous map ${}^\bullet f : {}^\bullet M \to {}^\bullet N$ is called *standard* if it is induced by a map $f : M \to N$ *i.e.* if for any ${}^\bullet x \in {}^\bullet M$ and any representative $\{x_\alpha\}$, $\{f(x_\alpha)\}$ is a representative of ${}^\bullet f({}^\bullet x)$.

Theorem 2. *Let L be a dense leaf of a riemannian lamination $\mathscr{L}$. Then for every leaf $L' \subset \mathscr{L}$,*

$$\widetilde{L'} \ \leq_{\mathsf{vir}} \ \widetilde{L}.$$

Proof. Fix a global metric d on $\mathscr{L}$ which agrees locally with the riemannian metric on the leaves. (By this we mean that in sufficiently small flow boxes, d agrees with the distance function of ρ in any plaque.) Let $\{\tilde{x}'_\alpha\} \subset \widetilde{L}'$ be any sequence, $\{x'_\alpha\}$ its projection to L'. Let $\{\tilde{x}_\alpha\} \subset \widetilde{L}$ be a sequence whose projection $\{x_\alpha\}$ to L is d-asymptotic to $\{x'_\alpha\}$. By transversal continuity of the metric, we deduce a sequence of K_α-quasiisometries, $K_\alpha \to 1$,

$$f_\alpha : D_\delta(x_\alpha) \to D_\delta(x'_\alpha),$$

for some $\delta > 0$, where $D_\delta(x)$ means the open ρ-ball of radius δ about x. Then if ${}^\bullet\tilde{x}$, ${}^\bullet\tilde{x}'$ are the virtual classes of $\{\tilde{x}_\alpha\}$, $\{\tilde{x}'_\alpha\}$, the sequence of quasiisometries $\{f_\alpha\}$ induces an isometry $D_\delta({}^\bullet\tilde{x}) \to D_\delta({}^\bullet\tilde{x}')$. Since L is dense, we may continue these isometries along geodesics to obtain a locally isometric surjection $U \to U'$, where U, U' are the galaxies containing ${}^\bullet\tilde{x}$, ${}^\bullet\tilde{x}'$. But since these spaces are simply connected, and the map is isometric, this surjection is a bijection. Hence it inverts to an isometry $U' \to U$. Repeating this for every ${}^\bullet$-class of sequence in $\widetilde{L}'$, we obtain the desired virtual subisometry $\widetilde{L}' \leq_{\mathsf{vir}} \widetilde{L}$. $\square$

Two riemannian manifolds have the same *virtual geometry* if their universal covers are virtually isometric.

Corollary 1. *Dense leaves of a riemannian lamination $\mathscr{L}$ have the same virtual geometry.*

3. THE FUNDAMENTAL GERM

Let $\mathscr{L}$ be a riemannian lamination, x a point contained in a transversal T, L a leaf accumulating at x and L_x the leaf containing x. Let $\mathsf{f} : T \to \mathbf{F}_*\mathscr{L}$ be a continuous section of the leaf-wise orthonormal frame bundle of $\mathscr{L}$ over T. Fix locally isometric universal covers $p : \widetilde{L} \to L$ and $p_x : \widetilde{L}_x \to L_x$. Denote $T_0 = T \cap L$, $\widetilde{T}_0 = p^{-1}(T_0)$ and let $\tilde{\mathsf{f}}_{\tilde{y}}$ denote the lift of f_y to a point $\tilde{y} \in \widetilde{T}_0$ covering y. We pick a basepoint $\tilde{x} \in \widetilde{L}_x$ lying over x with lifted frame $\tilde{\mathsf{f}}_{\tilde{x}}$.

Let $\tilde{y} \in \widetilde{T}_0$. For $r > 0$, the frames $\tilde{\mathsf{f}}_{\tilde{x}}$, $\tilde{\mathsf{f}}_{\tilde{y}}$ determine polar coordinates on the metric disks $D_r(\tilde{x})$, $D_r(\tilde{y})$. This yields in turn a canonical quasiisometry

$$f : D_r(\tilde{x}) \to D_r(\tilde{y})$$

given by the coordinate maps.

Let $\{x_\alpha\} \subset T_0$ be a sequence converging to x, $\{\tilde{x}_\alpha\} \subset \widetilde{T}_0$ any sequence covering $\{x_\alpha\}$. Then the frame sequence $\{\tilde{\mathsf{f}}_{\tilde{x}_\alpha}\}$ and the frame $\tilde{f}_{\tilde{x}}$ determine a sequence of K_α-quasiisometries

$$\Big\{ f_\alpha : D_{r_\alpha}(\tilde{x}) \longrightarrow D_{r_\alpha}(\tilde{x}_\alpha) \Big\}.$$

Since L accumulates at x, we may choose the sequence of radii $r_\alpha \to \infty$ so that $K_\alpha \to 1$. We deduce an isometry

$${}^\bullet f : \widetilde{L}_x \longrightarrow U \subset {}^\bullet\widetilde{L}$$

where U is the galaxy containing ${}^\bullet\tilde{x}$. The map ${}^\bullet f$ is called an f-*diophantine approximation* of x along L.

Definition 3. The **fundamental germ** of $\mathscr{L}$, based at x along L and f, is

$$[\![\pi]\!]_1(\mathscr{L}, L, x, \mathsf{f}) \;=\; \Big\{ {}^\bullet f \circ {}^\bullet g^{-1} \;\Big|\; {}^\bullet f, {}^\bullet g \text{ are } \mathsf{f}\text{-diophantine approximations of } x \text{ along } L \Big\}.$$

If $x \in L$, we shorten the notation to $[\![\pi]\!]_1(\mathscr{L}, x, \mathsf{f})$. The groupoid structure of $[\![\pi]\!]_1(\mathscr{L}, L, x, \mathsf{f})$ will be described in the next section.

Note 1. Suppose that $\mathscr{L}$ is a constant curvature riemannian foliation with dense leaf L modeled on the space form $\mathbb{M}^n = \mathbb{R}^n$ or $\mathbb{H}^n$. Then the frame field actually determines a sequence of uniquely defined *global* isometries $\{f_\alpha : \mathbb{M}^n \to \mathbb{M}^n\}$. Given G a group, nonstandard G is the group ${}^*G$ of all sequences $\{g_\alpha\} \subset G$ modulo the relation $\sim$ described in (1). Then an f-diophantine approximation is completely determined by the class ${}^*f \in {}^*\mathsf{Isom}(\mathbb{M}^n)$ of $\{f_\alpha\}$. We note that ${}^*\mathsf{Isom}(\mathbb{M}^n)$ is a subgroup of $\mathsf{Isom}({}^\bullet\mathbb{M}^n)$ (the group of isometries of ${}^\bullet\mathbb{M}^n$, *not* virtual isometries). Thus, if $\Gamma \cong \pi_1 L$ is the deck group of $\mathbb{M}^n \to L$, we have

$$ {}^*\Gamma \;\subset\; [\![\pi]\!]_1(\mathscr{L}, L, x, \mathsf{f}) \;\subset\; {}^*\mathsf{Isom}(\mathbb{M}^n). $$

The terminology f-diophantine approximation comes from the following example.

Example 1. Let $\mathscr{L}$ be the irrational foliation of the torus $\mathbb{T}^2$ by lines of slope $r \in \mathbb{R} \setminus \mathbb{Q}$. Define a representation $\rho : \mathbb{Z} \cong \pi_1\mathbb{S}^1 \to \mathsf{Homeo}(\mathbb{S}^1)$ by $\rho_n(\overline{y}) = \overline{y - nr}$, where $\overline{y}$ denotes the image of $y \in \mathbb{R}$ in $\mathbb{S}^1 = \mathbb{R}/\mathbb{Z}$. Then the suspension of ρ, $\mathscr{L}_\rho = (\mathbb{R} \times \mathbb{S}^1)/\mathbb{Z}$, is homeomorphic to $\mathscr{L}$. The map $\mathbb{R} \times \mathbb{S}^1 \to \mathbb{S}^1$ defined $(x, \overline{y}) \mapsto \overline{x}$ (*i.e.* the projection onto the first factor composed with the universal covering $\mathbb{R} \to \mathbb{S}^1$) induces a projection $\mathscr{L}_\rho \to \mathbb{S}^1$. Let $T \approx \mathbb{S}^1$ be a fiber of this projection passing through x. A frame section f along T is determined by an orientation of $\mathscr{L}$. In this case, an f-diophantine approximation of x is just a diophantine approximation of r. (Recall that a sequence $\{n_\alpha\} \subset \mathbb{Z}$ is called a diophantine approximation of $r \in \mathbb{R}$ if $\{\overline{rn_\alpha}\}$ converges to $\overline{0} \in \mathbb{S}^1$.) Thus if one denotes by ${}^*\mathbb{Z}_r$ the subgroup of ${}^*\mathbb{Z}$ consisting of classes of diophantine approximations of r, we obtain in agreement with the construction in [4], §4.4:

$$ [\![\pi]\!]_1(\mathscr{L}, L, x, \mathsf{f}) \;=\; {}^*\mathbb{Z}_r. $$

(Note: ${}^*\mathbb{Z}_r$ is an ideal if and only if r is rational.) If another frame field f' is used whose domain is a transversal T' which is not a suspension fiber, the set of diophantine approximations is a subset ${}^*\mathbb{R}_r \subset {}^*\mathbb{R}$. This subset maps injectively into ${}^\bullet\mathbb{R}$ with image ${}^\bullet\mathbb{Z}_r = {}^*\mathbb{Z}_r$.

Example 2. Consider a nested set of Fuchsian groups $\mathscr{G} = \{\Gamma_i\}$ and let

$$ \widehat{\Sigma}_{\mathscr{G}} \;=\; \varprojlim \mathbb{H}^2/\Gamma_i, $$

be the associated hyperbolic surface solenoid. We may take T to be a fiber $\hat{p}^{-1}(x_0)$ of the projection $\hat{p} : \widehat{\Sigma}_{\mathscr{G}} \to \Sigma_0$, where $\Sigma_0 = \mathbb{H}^2/\Gamma_0$ is the initial surface. Then a frame at x_0 pulls back to a frame section f along T. In this case, we find that Definition 3 again agrees with the definition found in [4]:

$$ \begin{aligned} [\![\pi]\!]_1(\widehat{\Sigma}_{\mathscr{G}}, L, x, \mathsf{f}) \quad &= \quad \bigcap {}^*\Gamma_i \\ &= \quad \left\{ \{g_\alpha\} \subset \Gamma_0 \;\middle|\; \text{for all } i,\ \exists\, N_i \text{ such that } g_\alpha \in \Gamma_i \text{ when } \alpha > N_i \right\} \Big/ \sim, \end{aligned} $$

a subgroup of ${}^*PSL(2,\mathbb{R}) \cong PSL(2, {}^*\mathbb{R})$. If f' is another frame field, not necessarily with a fiber transversal domain, then the corresponding germ $[\![\pi]\!]_1(\widehat{\Sigma}_{\mathscr{G}}, L, x, \mathsf{f}')$ need not define a subgroup of $PSL(2, {}^*\mathbb{R})$ and particularly, need not be isomorphic to $\bigcap {}^*\Gamma_i$ (although the fundamental germs calculated with respect to f and f' are in canonical bijection). The rub here is the non-uniform nature of the action of $PSL(2,\mathbb{R})$ on $\mathbb{H}^2$. This problem will become moot through the replacement of the fundamental germ by the mother germ, §6.

Example 3. More generally, let $\mathscr{L}$ be any hyperbolic surface lamination. Then the fundamental germ $[\![\pi]\!]_1(\mathscr{L}, L, x, \mathsf{f})$ is a subset of $PSL(2, {}^*\mathbb{R})$. Equally, if $\mathscr{L}$ is a hyperbolic 3-lamination, $[\![\pi]\!]_1(\mathscr{L}, L, x, \mathsf{f}) \subset PSL(2, {}^*\mathbb{C})$.

4. THE GERM UNIVERSAL COVER

Let $L \subset \mathscr{L}$ be a fixed leaf. Denote by $p : \widetilde{L} \to L$ the universal cover. We recall the following definition [4]:

Definition 4. The **germ universal cover** of $\mathscr{L}$ along L is the subspace $\llbracket \widetilde{\mathscr{L}} \rrbracket \subset {}^{\bullet}\widetilde{L}$ defined

$$\llbracket \widetilde{\mathscr{L}} \rrbracket \;=\; \left\{ \{\tilde{x}_\alpha\} \subset \widetilde{L} \;\middle|\; \{p(\tilde{x}_\alpha)\} \text{ converges in } \mathscr{L} \right\} \Big/ \simeq .$$

We will denote elements of the germ universal cover by ${}^{\bullet}\tilde{x}$. There is a natural projection

$${}^{\bullet}p : \llbracket \widetilde{\mathscr{L}} \rrbracket \longrightarrow \mathscr{L} \qquad\qquad {}^{\bullet}\tilde{x} \longmapsto \hat{x} \;=\; \lim p(\tilde{x}_\alpha),$$

where $\{\tilde{x}_\alpha\}$ is a representative sequence in the class ${}^{\bullet}\tilde{x}$. We will write $\lim {}^{\bullet}\tilde{x} = \hat{x}$ if ${}^{\bullet}p({}^{\bullet}\tilde{x}) = \hat{x}$. Note that ${}^{\bullet}p$ is surjective if and only if L is dense and in general ${}^{\bullet}p$ maps onto the closure $\overline{L}$ of L, itself a sublamination of $\mathscr{L}$.

Proposition 1. *$\llbracket \widetilde{\mathscr{L}} \rrbracket$ consists of a union of galaxies of ${}^{\bullet}\widetilde{L}$.*

Proof. Let ${}^{\bullet}\tilde{x} \in \llbracket \widetilde{\mathscr{L}} \rrbracket$ and denote by U the galaxy containing ${}^{\bullet}\tilde{x}$. If ${}^{\bullet}\tilde{y} \in U$, then there exists a sequence of geodesic paths $\{\tilde{\eta}_\alpha\}$ connecting representatives $\{\tilde{x}_\alpha\}$ to $\{\tilde{y}_\alpha\}$ in $\widetilde{L}$, whose projection to $\mathscr{L}$ gives a convergent sequence of paths $\{\eta_\alpha\}$. It follows that the projection $\{p(\tilde{y}_\alpha)\}$ converges, and ${}^{\bullet}\tilde{y} \in \llbracket \widetilde{\mathscr{L}} \rrbracket$ as well. $\square$

The galaxies that make up $\llbracket \widetilde{\mathscr{L}} \rrbracket$ will be referred to as leaves. See [4] for a proof of the following

Theorem 3. *$\llbracket \widetilde{\mathscr{L}} \rrbracket$ may be given the structure of a lamination whose leaves are nowhere dense and such that the map ${}^{\bullet}p : \llbracket \widetilde{\mathscr{L}} \rrbracket \to \overline{L}$ is an open surjection.*

One can thus think of $\llbracket \widetilde{\mathscr{L}} \rrbracket$ as a the result of unwrapping all of the diophantine approximations implied by L. The topology that $\llbracket \widetilde{\mathscr{L}} \rrbracket$ obtains from its lamination atlas is not unique, and is called a *germ universal cover topology*. It is in general coarser than the topology $\llbracket \widetilde{\mathscr{L}} \rrbracket$ induces from ${}^{\bullet}\widetilde{L}$.

Proposition 2. *If $\mathscr{L}$ is compact then $\llbracket \widetilde{\mathscr{L}} \rrbracket = {}^{\bullet}\widetilde{L}$.*

Proof. This follows from well-known compactness arguments *e.g.* see the proof in [4]. $\square$

An element ${}^{\bullet}u = {}^{\bullet}f \circ {}^{\bullet}g^{-1} \in \llbracket \pi \rrbracket_1(\mathscr{L}, L, x, \mathsf{f})$ arises as the limit of a sequence of K_α-quasiisometries

$$\left\{ u_\alpha : D_{r_\alpha}(\tilde{x}_\alpha) \longrightarrow D_{r_\alpha}(\tilde{y}_\alpha) \right\}, \tag{2}$$

where $\{\tilde{x}_\alpha\}, \{\tilde{y}_\alpha\} \subset L$, $K_\alpha \to 1$ and $r_\alpha \to \infty$. The limit ${}^{\bullet}u : V \to U$ is independent of the sequence $\{u_\alpha\}$ and depends only on the sequences of frames $\{\mathsf{f}_{x_\alpha}\}$, $\{\mathsf{f}_{y_\alpha}\}$. In particular, we could have obtained ${}^{\bullet}u$ through the same sequence of quasiisometries with domains extended to a sequence of larger disks $D_{s_\alpha}(\tilde{x}_\alpha)$, $s_\alpha > r_\alpha$ – provided that the new quasiisometry constants converge to 1 as well.

Now for arbitrary ${}^{\bullet}\tilde{x} \in \llbracket \widetilde{\mathscr{L}} \rrbracket$, the expression ${}^{\bullet}u({}^{\bullet}x)$ does not even make formal sense, since ${}^{\bullet}u$ is so far only defined on the galaxy V. We contrast this with the constant curvature case, where, because $\llbracket \pi \rrbracket_1(\mathscr{L}, x, L, \mathsf{f}) \subset \mathsf{Isom}({}^{\bullet}\mathbb{M})$, ${}^{\bullet}u({}^{\bullet}\tilde{x})$ is always formally defined, although it need not define an element of $\llbracket \widetilde{\mathscr{L}} \rrbracket$.

Let us say that ${}^\bullet u$ is *formally defined* on an element ${}^\bullet\tilde{w} \in [\![\widetilde{\mathscr{L}}]\!]$ if there exists a sequence (2) giving rise to ${}^\bullet u$ and a representative sequence $\{\tilde{w}_\alpha\}$ of ${}^\bullet\tilde{w}$ such that

$$D_{r'_\alpha}(w_\alpha) \subset D_{r_\alpha}(\tilde{x}_\alpha)$$

for all α, where $r'_\alpha \to \infty$. It follows then that if V' is the galaxy containing ${}^\bullet\tilde{w}$, then the limit ${}^\bullet u$ is defined on V' as well. Whenever we write ${}^\bullet u({}^\bullet w)$, it will tacitly be understood that ${}^\bullet u$ is formally defined at ${}^\bullet w$.

Define the domain of ${}^\bullet u$ as

$$\mathsf{Dom}({}^\bullet u) \;=\; \left\{ {}^\bullet\tilde{x} \in [\![\widetilde{\mathscr{L}}]\!] \;\middle|\; {}^\bullet u({}^\bullet\tilde{x}) \in [\![\widetilde{\mathscr{L}}]\!] \text{ and } \lim {}^\bullet u({}^\bullet\tilde{x}) = \lim {}^\bullet\tilde{x} \right\},$$

and $\mathsf{Ran}({}^\bullet u) = {}^\bullet u(\mathsf{Dom}({}^\bullet u))$. With this definition, it follows that $[\![\pi]\!]_1(\mathscr{L}, L, x, \mathsf{f})$ has the structure of a groupoid. Note that for any ${}^\bullet u \in [\![\pi]\!]_1(\mathscr{L}, L, x, \mathsf{f})$, $\mathsf{Dom}({}^\bullet u)$, $\mathsf{Ran}({}^\bullet u)$ are unions of leaves and hence induce lamination structures from $[\![\widetilde{\mathscr{L}}]\!]$. Moreover, on $\mathsf{Dom}({}^\bullet u)$,

$$ {}^\bullet p \circ {}^\bullet u \;=\; {}^\bullet p. \tag{3}$$

In particular we see that ${}^\bullet u : \mathsf{Dom}({}^\bullet u) \to \mathsf{Ran}({}^\bullet u)$ defines a lamination homeomorphism.

Example 4. Let $\mathscr{L}$ be the irrational foliation of $\mathbb{T}^2$ by lines of slope r, $L \approx \mathbb{R}$ any dense leaf. Then by Proposition 2, $[\![\widetilde{\mathscr{L}}]\!] = {}^\bullet\mathbb{R}$. Moreover, for any frame field f and ${}^\bullet u \in [\![\pi]\!]_1(\mathscr{L}, L, x, \mathsf{f})$, it is not difficult to see that $\mathsf{Dom}({}^\bullet u) = {}^\bullet\mathbb{R}$. Thus, $[\![\pi]\!]_1(\mathscr{L}, L, x, \mathsf{f})$ is a group isomorphic to ${}^*\mathbb{Z}_r$.

Example 5. Let $\mathscr{L}$ be the profinite hyperbolic surface solenoid $\widehat{\Sigma}_{\mathscr{G}}$ of *Example* 2. Then we have, again by compactness,

$$[\![\widetilde{\mathscr{L}}]\!] \;=\; {}^\bullet\mathbb{H}^2.$$

If f is a frame field lifted from a frame on a surface occurring in the defining inverse limit, then $[\![\pi]\!]_1(\mathscr{L}, L, x, \mathsf{f})$ is a group. On the other hand, if f is a frame field not obtained in this way, then $[\![\pi]\!]_1(\mathscr{L}, L, x, \mathsf{f})$ need not be a group *e.g.* see §6.

The following may also be found in [4].

Theorem 4. *Let $F : (\mathscr{L}, L) \to (\mathscr{L}', L')$ be a lamination map. Then there exist germ universal cover topologies so that the map*

$$[\![\widetilde{F}]\!] : [\![\widetilde{\mathscr{L}}]\!] \longrightarrow [\![\widetilde{\mathscr{L}'}]\!]$$

induced by $\{\tilde{x}_\alpha\} \mapsto \{\widetilde{F}(\tilde{x}_\alpha)\}$ is a continuous lamination map.

Note 2. It is useful here to point out that for a lamination $\mathscr{L}_\rho = (\widetilde{B} \times \mathsf{F})/\pi_1 B$ occurring as a suspension of a representation $\rho : \pi_1 B \to \mathsf{Homeo}(\mathsf{F})$, it is in general false that a lamination homeomorphism $F : \mathscr{L}_\rho \to \mathscr{L}_\rho$ lifts to a homeomorphism of the "universal covering space" $\widetilde{B} \times \mathsf{F}$.

Now suppose that L' is another leaf of $\mathscr{L}$. Denote by $[\![\widetilde{\mathscr{L}}]\!]'$ the germ universal cover formed from L'.

Proposition 3. *If L' accumulates on L then there is a virtual subisometry ${}^\bullet\widetilde{L} \to {}^\bullet\widetilde{L}'$ restricting to a virtual subisometry*

$$[\![\widetilde{\mathscr{L}}]\!] \;\longrightarrow\; [\![\widetilde{\mathscr{L}}]\!]'$$

which is a homeomorphism onto its image with respect to appropriate germ universal cover topologies.

Proof. This follows directly from the proof of Theorem 2. □

5. SENSITIVITY TO CHANGES IN DATA

In this section we shall examine the dependence of the fundamental germ on the base point x, the accumulating leaf L and the frame field f.

Change in base point and accumulating leaf: Let us fix for the moment the dense leaf L and consider a change of base point $x \mapsto x'$ in which $L_x = L_{x'}$. Let η be a geodesic connecting x to x' in L_x. The tangent vector v to η at x has coordinate $(a_1, \dots, a_n)$ with respect to the frame f_x. At each $y \in T =$ the domain of f, this coordinate determines a vector v_y using the frame f_y. We obtain in this way a transversally continuous family of geodesics $\{\eta_y\}_{y \in T}$. Restricting to an open subtransversal of T if necessary, we may parallel translate f along the geodesic family to obtain a frame field f' with domain $T' \ni x'$. The following is then immediate from the definition of the fundamental germ.

Proposition 4. *Let x', f' be as in the preceding paragraph. Then*

$$[\![\pi]\!]_1(\mathscr{L}, L, x, \mathsf{f}) \;=\; [\![\pi]\!]_1(\mathscr{L}, L, x', \mathsf{f}').$$

If we consider a change of base point $x \mapsto x'$, in which $L_x \neq L_{x'}$, the situation becomes considerably more subtle. In fact, we shall see in §7 that fundamental germs based at points on different leaves can be nonisomorphic. For similar reasons, a change in accumulating leaf L may yield nonisomorphic fundamental germs.

Change in frame field: Let us now fix the base point x and consider a new frame field $\mathsf{f}' : T' \to \mathbf{F}_*\mathscr{L}$ based at x. For simplicity, we again assume that $\pi_1 L = 1$. Since T (the domain of f) and T' each contain subtransversal neighborhoods of x lying in a common flow box, it is clear that there is a natural bijection

$$[\![\pi]\!]_1(\mathscr{L}, L, x, \mathsf{f}) \;\longleftrightarrow\; [\![\pi]\!]_1(\mathscr{L}, L, x, \mathsf{f}').$$

The issue is then the law of composition. We will show that this map need not be an isomorphism.

Let us consider the inverse limit solenoid $\widehat{\Sigma}_{\mathscr{G}}$ of *Example* 2. Assume that $L = L_x$, $T = T' =$ a fiber over a point $x_0 \in \Sigma_0$ and that $\mathsf{f}_x = \mathsf{f}'_x$. We will take f to be simply the lift of a frame based at x_0, so that f-diophantine approximations consist of sequences $\{\gamma_\alpha\} \subset \Gamma_0$ converging with respect to the family $\{\Gamma_i\}$. It follows that every f'-diophantine approximation of x may be written in the form $\{\gamma_\alpha \Theta_\alpha\}$, where $\{\Theta_\alpha\}$ consists of a sequence of rotations based at x with angle going to 0 and $\{\gamma_\alpha\}$ is an f-diophantine approximation. General elements of $[\![\pi]\!]_1(\mathscr{L}, L, x, \mathsf{f}')$ are then of the form $\{\gamma_\alpha \Theta_\alpha \eta_\alpha\}$, where $\{\eta_\alpha\}$ is another f-diophantine approximation. We should not expect products of elements of this type to yield elements of $[\![\pi]\!]_1(\mathscr{L}, L, x, \mathsf{f}')$. Indeed, such a product would have the shape

$$\{\gamma_\alpha \Theta_\alpha \eta_\alpha \Delta_\alpha \omega_\alpha\}, \tag{4}$$

for $\{\Delta_\alpha\}$ another sequence of rotations based at x with angle going to 0 and $\{\omega_\alpha\}$ another f-diophantine approximation. If Θ_α does not converge to the identity fast enough, $\Theta_\alpha \eta_\alpha \Delta_\alpha \omega_\alpha$ applied to f'_x will not project to a frame based at $x_0 \in \Sigma_0$. Hence the expression (4) is not even asymptotic to an element of $[\![\pi]\!]_1(\mathscr{L}, L, x, \mathsf{f}')$. It is not difficult to see that unless f' is the pull-back of a frame on Σ_0, this sort of problem always arises.

6. THE MOTHER GERM

In this section, we assume that $\mathscr{L}$ has a dense leaf L, with which we define the germ universal cover $[\![\widetilde{\mathscr{L}}]\!]$, equipped with a fixed germ universal cover topology.

While the fundamental germ $[\![\pi]\!]_1(\mathscr{L},L,x,\mathsf{f})$ enjoys the property of being reasonably calculable and leaf specific, it can be sensitive to data variation. There are additional shortcomings:

- By (3), the action of the fundamental germ $[\![\pi]\!]_1(\mathscr{L},L,x,\mathsf{f})$ on $[\![\widetilde{\mathscr{L}}]\!]$ respects the germ covering ${}^{\bullet}p$. However it need not be the case that every identification implied by ${}^{\bullet}p$ is implemented by an element of $[\![\pi]\!]_1(\mathscr{L},L,x,\mathsf{f})$.
- There will be in general other maps of leaves of $[\![\widetilde{\mathscr{L}}]\!]$ that satisfy (3) but do not appear in $[\![\pi]\!]_1(\mathscr{L},L,x,\mathsf{f})$.
- It appears that $[\![\pi]\!]_1(\mathscr{L},L,x,\mathsf{f})$ such as it is defined, will be functorial only under certain types of lamination maps *e.g.* see [4].

For this reason, we will expand $[\![\pi]\!]_1(\mathscr{L},L,x,\mathsf{f})$ to a larger groupoid, called the mother germ. The mother germ will be the maximal amplification of $[\![\pi]\!]_1(\mathscr{L},L,x,\mathsf{f})$ which contains all partially defined maps of sublaminations of $[\![\widetilde{\mathscr{L}}]\!]$ satisfying (3): in other words, it is the full deck groupoid of ${}^{\bullet}p$.

Let $\mathsf{Dom}, \mathsf{Ran}$ be sublaminations of $[\![\widetilde{\mathscr{L}}]\!]$. A homeomorphism

$$ {}^{\bullet}u : \mathsf{Dom} \longrightarrow \mathsf{Ran} \ \subset\ [\![\widetilde{\mathscr{L}}]\!] $$

satisfying (3) is called *deck*. Note that condition (3) implies that a deck homeomorphism ${}^{\bullet}u$ is automatically an isometry along the leaves of Dom.

Definition 5. The **mother germ** is the groupoid

$$ [\![\pi]\!]_1(\mathscr{L}) \ = \ \Big\{ {}^{\bullet}u : \mathsf{Dom} \longrightarrow \mathsf{Ran}\text{ is a deck homeomorphism} \Big\}. $$

The mother germ will never be a group, since it distinguishes deck maps obtained from others by restriction of domain. In general, however, it will contain many interesting and calculable subgroups and subgroupoids, as the following shows.

Proposition 5. *Let L' be any dense leaf of $\mathscr{L}$. Then there is an injective groupoid homomorphism*

$$ [\![\pi]\!]_1(\mathscr{L},L',x,\mathsf{f}) \ \hookrightarrow\ [\![\pi]\!]_1(\mathscr{L}). $$

Proof. By Proposition 3, there exists an isometric inclusion ${}^{\bullet}f : [\![\widetilde{\mathscr{L}}]\!]' \hookrightarrow [\![\widetilde{\mathscr{L}}]\!]$. If ${}^{\bullet}u \in [\![\pi]\!]_1(\mathscr{L},L',x,\mathsf{f})$, then the map

$$ {}^{\bullet}u \ \longmapsto\ {}^{\bullet}f \circ {}^{\bullet}u \circ {}^{\bullet}f^{-1} $$

defines an injective groupoid homomorphism. ☐

Theorem 5. *The quotient*

$$ [\![\pi]\!]_1(\mathscr{L})\backslash[\![\widetilde{\mathscr{L}}]\!], $$

equipped with the quotient germ universal cover topology, has the structure of a riemannian lamination canonically isometric to $\mathscr{L}$.

Proof. Let ${}^{\bullet}\tilde{x}$, ${}^{\bullet}\tilde{y} \in [\![\widetilde{\mathscr{L}}]\!]$ be such that $\lim {}^{\bullet}\tilde{x} = \lim {}^{\bullet}\tilde{y} = x$. Thus each point is represented by sequences in $\widetilde{L}$ that project to sequences $\{x_\alpha\}$, $\{y_\alpha\} \subset L$ having a common limit x. Let f be a frame field along a transversal T containing x and which we may assume contains $\{x_\alpha\}$ and $\{y_\alpha\}$. Then if ${}^{\bullet}f$, ${}^{\bullet}g$ are the diophantine approximations associated to $\{x_\alpha\}$, $\{y_\alpha\}$ we have ${}^{\bullet}u = {}^{\bullet}g \circ {}^{\bullet}f^{-1} \in [\![\pi]\!]_1(\mathscr{L},L,x,\mathsf{f})$ identifies ${}^{\bullet}\tilde{x}$ with ${}^{\bullet}\tilde{y}$. Since this latter groupoid belongs to the mother germ by Proposition 5, it follows that $[\![\pi]\!]_1(\mathscr{L})\backslash[\![\widetilde{\mathscr{L}}]\!]$ contains all of the identifications implied by ${}^{\bullet}p$ and so may be identified with $\mathscr{L}$ with its quotient topology. ☐

Let $\mathscr{L}$, $\mathscr{L}'$ be riemannian laminations with dense leaves L, L'. A map $[\![\widetilde{F}]\!] : [\![\widetilde{\mathscr{L}}]\!] \to [\![\widetilde{\mathscr{L}'}]\!]$ is called *standard* if it is induced by $F : \widetilde{L} \to \widetilde{L}'$ (*e.g.* compare with the definition found in §3). In addition, $[\![\widetilde{F}]\!]$ is called $[\![\pi]\!]_1(\mathscr{L})$*-equivariant* if there exists a groupoid homomorphism $[\![F]\!]_* : [\![\pi]\!]_1(\mathscr{L}) \to [\![\pi]\!]_1(\mathscr{L}')$ such that

$$[\![\widetilde{F}]\!]\big({}^{\bullet}u \cdot {}^{\bullet}\tilde{x}\big) \;=\; [\![F]\!]_*\big({}^{\bullet}u\big) \cdot [\![\widetilde{F}]\!]\big({}^{\bullet}\tilde{x}\big)$$

for all ${}^{\bullet}u \in [\![\pi]\!]_1(\mathscr{L})$ and ${}^{\bullet}\tilde{x} \in \mathsf{Dom}({}^{\bullet}u)$.

Theorem 6. *Let $[\![\widetilde{F}]\!] : [\![\widetilde{\mathscr{L}}]\!] \to [\![\widetilde{\mathscr{L}'}]\!]$ be a standard, $[\![\pi]\!]_1(\mathscr{L})$-equivariant map. Then $[\![\widetilde{F}]\!]$ covers a unique map $F : \mathscr{L} \to \mathscr{L}'$.*

Proof. By equivariance, the expression

$$F \;=\; {}^{\bullet}p' \circ [\![\widetilde{F}]\!] \circ {}^{\bullet}p^{-1}$$

yields a well-defined function $F : \mathscr{L} \to \mathscr{L}'$, continuous because ${}^{\bullet}p, {}^{\bullet}p'$ are open maps and $[\![\widetilde{F}]\!]$ is continuous. ☐

In [4], functoriality of the fundamental germ was demonstrated only with respect to the restricted class of *trained* lamination maps. The following theorem shows that the mother germ is considerably more flexable. A *lamination covering map* is a surjective lamination map which is a covering map when restricted to any leaf.

Theorem 7. *Let $F : \mathscr{L} \to \mathscr{L}'$ be a lamination covering map. Then F induces an injective homomorphism of mother germs*

$$[\![F]\!]_* : [\![\pi]\!]_1(\mathscr{L}) \;\hookrightarrow\; [\![\pi]\!]_1(\mathscr{L}').$$

Proof. Let $L \subset \mathscr{L}$ be a dense leaf and let $\widetilde{F} : \widetilde{L} \to \widetilde{L}'$ be the leaf universal cover lift. Then by Theorem 4, $\widetilde{F}$ induces a standard map

$$[\![\widetilde{F}]\!] : [\![\widetilde{\mathscr{L}}]\!] \longrightarrow [\![\widetilde{\mathscr{L}'}]\!].$$

We note that since $\widetilde{F}$ is injective, $[\![\widetilde{F}]\!]$ is a homeomorphism onto its image lamination. Let ${}^{\bullet}u \in [\![\pi]\!]_1(\mathscr{L})$. Then the map

$$[\![F]\!]_*\big({}^{\bullet}u\big) \;:=\; [\![\widetilde{F}]\!] \circ {}^{\bullet}u \circ [\![\widetilde{F}]\!]^{-1},$$

defined on $[\![\widetilde{F}]\!]\big(\mathsf{Dom}({}^{\bullet}u)\big)$, is deck for the germ universal covering ${}^{\bullet}p'$. Indeed

$$\begin{aligned} {}^{\bullet}p' \circ \Big([\![\widetilde{F}]\!] \circ {}^{\bullet}u \circ [\![\widetilde{F}]\!]^{-1}\Big) &= F \circ \big({}^{\bullet}p \circ {}^{\bullet}u\big) \circ [\![\widetilde{F}]\!]^{-1} \\ &= F \circ {}^{\bullet}p \circ [\![\widetilde{F}]\!]^{-1} \\ &= {}^{\bullet}p'. \end{aligned}$$

Thus the map $[\![F]\!]_*$ is an injective groupoid homomorphism, and we are done. ☐

We have the following

Corollary 2. *The mother germ $[\![\pi]\!]_1(\mathscr{L})$ is independent of leaf-wise riemannian metric and smooth structure. In particular, $[\![\pi]\!]_1(\mathscr{L})$ is a* topological *invariant.*

7. THE ANTENNA LAMINATION

In this section, we will calculate the fundamental germ of the antenna Riemann surface lamination of Kenyon and Ghys [6]: it is distinguished by the unusual property of having dense leaves of both planar and hyperbolic conformal type.

We begin by constructing a graphical model of a dense leaf of the antenna lamination. Let T_1 be the cross with vertices $V_1 = \big\{(0,0),(\pm 1,0),(0,\pm 2)\big\}$ and edges consisting of the line segments connecting $(0,0)$ to each of the other four vertices. Suppose that we have constructed T_n meeting the x-axis in the interval $[-2^n+1, 2^n-1]\times\{0\}$ and meeting the y-axis in the interval $\{0\}\times[-2^n,2^n]$. Translate T_n vertically so that the origin is taken to $(0,2^n)$ and consider the images of this translate by rotations of the plane – about the origin – of angles $0, \pm\pi/2, \pi$. The union of these images forms a tree; T_{n+1} is then obtained by replacing the extremal edges $[2^{n+1}-2, 2^{n+1}]\times\{0\}$ and $[-2^{n+1}, -2^{n+1}+2]\times\{0\}$ by $[2^{n+1}-2, 2^{n+1}-1]\times\{0\}$ and $[-2^{n+1}+1, -2^{n+1}+2]\times\{0\}$. It follows that $T_1\subset T_2\subset\dots$: we then define

$$T_\infty \;=\; \varinjlim T_n.$$

See Figure 1.



FIGURE 1. The Antenna Tree

Given $n\in\mathbb{Z}$, let $\mathsf{ord}_2(n)$ be the 2-adic order: the largest nonnegative integer r for which 2^r divides n. Then the vertex set of T_∞ is

$$V_\infty \;=\; \big\{(0,0)\big\}\bigcup\Big\{v=(x,y)\in\mathbb{Z}\oplus\mathbb{Z}\;\Big|\;\mathsf{ord}_2(x)\neq\mathsf{ord}_2(y)\Big\}.$$

We may view V_∞ as a groupoid through its action on itself by addition. In order to avoid confusion, we write $v\circ w$ to indicate groupoid composition, in order to distinguish it from the element $v+w\in\mathbb{Z}\oplus\mathbb{Z}$.

Proposition 6. *For all $v,w\in V_\infty$, the composition $v\circ w$ is defined if and only if $v=-w$.*

Proof. Let $v,w\in V_\infty$. We show that $\mathsf{Ran}(w)=\mathsf{Dom}(v)$ if and only if $v=-w$. Suppose $v\neq -w$. Then we may write

$$v+w \;=\; \left(\sum_{\alpha=m}^{M} a_\alpha 2^\alpha, \sum_{\alpha=n}^{N} b_\alpha 2^\alpha\right)$$

where m, n are the first non-zero indices of the 2-adic expansions of the coordinates. If $v \circ w$ is defined, then since $0 \in \mathsf{Dom}(w)$, we must have $v + w \in V_\infty$. In particular, at least one of m or n is nonzero. Suppose it is n; we may assume without loss of generality that $m < n$. Write

$$w = \left(\sum_{\alpha=r}^{R} c_\alpha 2^\alpha, \sum_{\alpha=s}^{S} d_\alpha 2^\alpha \right).$$

Let $x = (0, 2^m)$. If $r < m$, then $x \in \mathsf{Dom}(w)$ but $v + w + x \notin V_\infty$ *i.e.* $\mathsf{Ran}(w) \neq \mathsf{Dom}(v)$. This is also true when $r \geq m$ except for two cases. If $r > m$ and $s = m$, $w + x$ is not defined presisely when $1 = d_s = \cdots = d_{r-1}$ and $d_r = 0$. Here we take $x' = (2^r, 2^m) \in \mathsf{Dom}(w)$ and note that $v + w + x' \notin V_\infty$. If $r = m$ and $s > m$, then $w + x$ is not defined. In this case, it follows from the form of $v + w$ that if $v = (v_1, v_2)$ then $\mathsf{ord}_2(v_1) > m$, so that $x \in \mathsf{Dom}(v)$. On the other hand, $x - w \notin \mathsf{Dom}(w)$. Thus $\mathsf{Ran}(w) \neq \mathsf{Dom}(v)$ here as well. $\square$

The lines $x = \pm y$ intersect T_∞ at the origin only. Each of the four components of $T_\infty \setminus \{(0,0)\}$ defines an end, one contained in each of the four components of $\mathbb{R}^2 \setminus \{(x, \pm x) \mid x \in \mathbb{R}\}$. Equipped with the path metric induced from $\mathbb{R}^2$, T_∞ has exactly four orientation preserving isometries, corresponding to the rotations about the origin of angles $0, \pm\pi/2, \pi$ (since ends must be taken to ends). On the other hand, T_∞ has many partially defined isometries. For example, for $v \in V_\infty$, let I_v be the map of $\mathbb{Z} \oplus \mathbb{Z}$ defined

$$I_v(x,y) = v + (x,y).$$

Then there is a maximal subtree $T_\infty^v \subset T_\infty$ (not necessarily connected) for which $I_v(T_\infty^v) \subset T_\infty$. By definition, I_v is isometric on its domain of definition. If v has coordinates of large 2-adic order, then I_v is defined on a large ball about 0 in T_∞. More precisely, if $v = (x,y)$ and $\mathsf{ord}_2(x), \mathsf{ord}_2(y) \geq n$ then $T_n \subset T_\infty^v$. Although the inverse $I_v^{-1} = I_{-v}$ is always defined at 0, the composition $I_{v_1} \circ I_{v_2} = I_{v_1+v_2}$ will not be defined at 0 if $v_1 + v_2 \notin V_\infty$.

We now define a riemannian surface modelled on T_∞, which will occur as a dense leaf of the antenna lamination. Regarding $T_\infty \subset \mathbb{R}^2 \times \{0\} \subset \mathbb{R}^3$, it is clear that

$$S_\infty = \text{boundary of a tubular neighborhood of } T_\infty$$

is homeomorphic to a sphere with four punctures. We want to fix a particular realization of S_∞ so that the partial isometries I_v of T_∞ will induce partial isometries of S_∞. Torward this end, consider the surfaces shown in Figure 2. We assume that they are equipped with



FIGURE 2. Building Blocks

riemannian metrics and boundary parametrizations so that given any pair of such surfaces and a choice of boundary component of each, the glueings are canonical and isometric. Each riemannian surface corresponds to a subgraph of T_∞, and we may build S_∞ from these riemannian surfaces using T_∞ as a template. The metrics on the building blocks will also be chosen so that when S_∞ is assembled within $\mathbb{R}^3$ it is invariant not only with respect to $\pi/2$-rotations about the z-axis, but also π-rotations about the x and y-axes. We think of T_∞ as a spine floating inside the tubular neighborhood bounded by S_∞, and we project in the positive vertical direction a copy of T_∞ onto S_∞. We denote this copy also by T_∞, and use the symbols 0 and v to denote the origin and a generic element of its vertex set as well. Having constructed S_∞ in this way, it is clear that every I_v induces a partial isometry of S_∞ whose domain is the subsurface (with boundary) of S_∞ modelled on T^v_∞. We denote this partial isometry I_v as well.

Let S^+_∞ be the intersection of S_∞ with the half plane $z \geq 0$. The universal cover $\widetilde{S}_\infty$ of S_∞ is built up from "tiles" modelled on S^+_∞, glued together side by side according to the same pattern one uses to glue ideal quadrilaterals to obtain the hyperbolic plane as the universal cover of the four times punctured sphere. Fix $\tilde{0} \in \widetilde{S}_\infty$ a base point lying over 0. The deck group of the universal covering map is F_3, the free group on three generators.

Let w_0 be the unit vector based 0 which is parallel to the x-axis and points in the positive direction. Consider the vector field W on the vertices of T_∞ obtained by parallel translating w_0 along T_∞. Note that the partial isometry of S_∞ induced by I_v, $v \in V_\infty$, takes w_0 to $\mathsf{w}_v = \mathsf{W}(v)$. This is not true of the rotations by angles $\pm\pi/2$ and π.

Let $\mathsf{D}_{\tilde{0}} \subset \widetilde{S}_\infty$ be the fundamental domain containing $\tilde{0}$. We lift T_∞ to $\mathsf{D}_{\tilde{0}}$, then translate it by F_3 to obtain a (disconnected) graph $\widetilde{T}_\infty$ on $\widetilde{S}_\infty$. Let $\widetilde{\mathsf{W}}$ be the vector field defined on the vertices of $\widetilde{T}_\infty$ that is the lift of W. The partial isometry I_v lifts to a partial isometry of $\widetilde{S}_\infty$ which maps a region of each fundamental domain D into D: we denote this privileged lift by I_v as well, and the set of such privileged lifts is denoted $\mathbb{I}$. In addition, by composing with elements of F_3, we obtain new partial isometries covering $I_v : S_\infty \to S_\infty$. We denote by $\widetilde{\mathbb{I}}$ the set of partial isometries of $\widetilde{S}_\infty$ obtained in this way. Then $F_3, \mathbb{I} \subset \widetilde{\mathbb{I}}$, and every element of F_3 commutes with every element of $\mathbb{I}$.

We are now ready to describe the antenna lamination. Consider first the space $\mathbb{A}$ of all trees in $\mathbb{R}^2$ whose vertex set contains the origin 0 and lies within $\mathbb{Z} \oplus \mathbb{Z}$. Each tree $T \in \mathbb{A}$ is equipped with the path metric induced from $\mathbb{R}^2$. On $\mathbb{A}$, we consider the metric

$$d(T,T') \;=\; \exp(-n),$$

where n is the largest integer such that the ball of radius n about 0 in T coincides with that about 0 in T'. $\mathbb{A}$ is a compact metric space, [6]. Two graphs T and T' are termed equivalent if there exists a translation by $(x,y) \in \mathbb{Z} \oplus \mathbb{Z}$ such that $T + (x,y) = T'$.

Now for any tree $T \in \mathbb{A}$, the ball of radius 1 about 0 is a tree $P \in \mathbb{A}$ all of whose vertices lie in the set $\{(0,0)\} \cup \{(\pm 1, \pm 1)\}$. We write $|P| \leq 4$ for the number of vertices v of P different from 0. There are 16 possible such P, and we may decompose $\mathbb{A}$ into a disjoint union of clopens $\mathbb{A}_P$, where $\mathbb{A}_P$ consists of those trees whose unit ball about 0 is P.

For each P, we consider in the spirit of Figure 2 a model pointed Riemann surface (Σ_P, z_P) homeomorphic to $\mathbb{S}^2 \setminus (|P|$ open disks$)$. We assume as before that each boundary component $\partial_v \Sigma_P$ – labeled by a vertex $v \neq (0,0)$ of P – has a fixed paramentrization, so that any two may be identified along their boundaries without ambiguity. Define

$$\mathscr{L} \;=\; \Big(\bigcup \left(\mathbb{A}_P \times \Sigma_P\right)\Big)\Big/ \text{gluing},$$

where the gluing is performed as follows. Given $T \in \mathbb{A}_P$, $v \in P$, the translate $T+v$ is in $\mathbb{A}_{P'}$ for some P', where $-v \in P'$. We then glue the boundaries $\partial_v \Sigma_P$ and $\partial_{-v}\Sigma_{P'}$. These gluings are compatible with the trivial lamination structures on the $\mathbb{A}_P \times \Sigma_P$ and thus $\mathscr{L}$ has the structure of a riemannian surface lamination. Note that there is an embedding $\mathbb{A} \hookrightarrow \mathscr{L}$ induced by $\mathbb{A}_P \times \{z_P\} \hookrightarrow \mathbb{A}_P \times \Sigma_P$.

Each leaf $L \subset \mathscr{L}$ corresponds to an equivalence class of graph $T \in \mathbb{A}$, embedded in L as a spine. Note that S_∞ is the leaf corresponding to the class of T_∞. Define the *antenna lamination* $\mathscr{L}_\infty$ to be the closure of S_∞ in $\mathscr{L}$.

Denote by $S_n \subset S_\infty$ the surface (with boundary) modelled on the subgraph $T_n \subset T_\infty$. If centered at a vertex $v \in V_\infty$ there is a subgraph isometric to T_n, it models a subsurface $S_n(v)$ containing v, and the isometry I_v maps S_n to $S_n(v)$.

The closure of V_∞ in $\mathscr{L}_\infty$ defines a transversal T through $0 \in S_\infty$, and the vector field W is transversally continuous with respect to the topology of T. A point $v \in V_\infty$ is transversally close to 0 if and only if its coordinates have large 2-adic order.

We are now ready to calculate the fundamental germ

$$[\![\pi]\!]_1(\mathscr{L}_\infty, 0, \mathsf{f}),$$

where f is the orthonormal frame field determined by W.

Define nested sets

$$\widetilde{G} \;=\; \{\widetilde{G}_n\} \;\subset\; \widetilde{\mathbb{I}} \quad \text{and} \quad G \;=\; \{G_n\} \;\subset\; \mathbb{I},$$

$n = 0, 1, 2, \ldots$, as follows. We say that $\widetilde{I} \in \widetilde{\mathbb{I}}$ is *n-close* if the domain of $\widetilde{I}$ contains the finite tree $\widetilde{T}_n \subset \widetilde{T}_\infty \cap \mathsf{D}_{\tilde{0}}$ corresponding to T_n, and maps it into the fundamental domain containing $\widetilde{I}(\tilde{0})$. Then $\widetilde{G}_n$ consists of the set of n-close maps and G_n the n-close maps in $\mathbb{I}$. Observe that

$$F_3 \;=\; \bigcap \widetilde{G}_n.$$

For $I \in G_n$, $I^{-1} \in G_n$ also. Moreover, if $I' \in G_m$ and the composition $I \circ I'$ is defined at $\tilde{0}$, then it belongs to G_N, for $N = \min(m, n)$.

Let

$$[\![G]\!] = \left\{ \{I_{v_\alpha} \circ I^{-1}_{v'_\alpha}\} \;=\; \{I_{v_\alpha - v'_\alpha}\} \;\middle|\; \{I_{v_\alpha}\},\, \{I_{v'_\alpha}\} \subset \mathbb{I} \text{ and converge w.r.t } G \right\} \Big/ \sim,$$

where the relation $\sim$ is defined by an ultrafilter $\mathfrak{U}$ as in (1). We denote the elements of $[\![G]\!]$ by $I_{{}^*v}$ where ${}^*v \in {}^*\mathbb{Z} \oplus {}^*\mathbb{Z}$, and regard $[\![G]\!]$ as a groupoid with unit space $[\![\widetilde{\mathscr{L}_\infty}]\!]$, in which the domains of elements are taken to be maximal in the sense defined in §4.

Proposition 7. *As a set,* $[\![G]\!]$ *may be identified with* ${}^*\mathbb{Z}_{\hat{2}} \oplus {}^*\mathbb{Z}_{\hat{2}}$, *where*

$${}^*\mathbb{Z}_{\hat{2}} \;=\; \left\{ \{n_\alpha\} \subset \mathbb{Z} \;\middle|\; \mathsf{ord}_2(n_\alpha) \to \infty \textit{ as } \alpha \to \infty \right\} \Big/ \sim .$$

Given ${}^*v, {}^*w \in {}^*\mathbb{Z}_{\hat{2}} \oplus {}^*\mathbb{Z}_{\hat{2}}$, *the composition* $I_{{}^*v} \circ I_{{}^*w}$ *is defined if and only if* ${}^*v = -{}^*w$.

Proof. Any element $({}^*n_1, {}^*n_2) \in {}^*\mathbb{Z}_{\hat{2}} \oplus {}^*\mathbb{Z}_{\hat{2}}$ may be written $({}^*n_1, 0) - (0, -{}^*n_2)$ which clearly defines an element of $[\![G]\!]$. Now consider ${}^*v, {}^*w \in {}^*\mathbb{Z}_{\hat{2}} \oplus {}^*\mathbb{Z}_{\hat{2}}$, and suppose that ${}^*v \neq -{}^*w$. The 2-adic order extends to

$$\mathsf{ord}_2 : {}^*\mathbb{Z}_{\hat{2}} \to {}^*\mathbb{Z}_\infty = ({}^*\mathbb{Z} \setminus \mathbb{Z}) \cup \{0\}.$$

Let ${}^*V_\infty \subset {}^*\mathbb{Z}_{\hat{2}} \oplus {}^*\mathbb{Z}_{\hat{2}}$ be the subset of pairs ${}^*u = ({}^*u_1, {}^*u_2)$ for which $\mathsf{ord}_2({}^*u_1) \neq \mathsf{ord}_2({}^*u_2)$. We distinguish four cases depending on whether ${}^*v, {}^*w \in {}^*V_\infty$ or not. If ${}^*v, {}^*w \in {}^*V_\infty$ then we may regard each as a class of sequences $\{v_\alpha\}$, $\{w_\alpha\} \subset V_\infty$. If $\{m_\alpha\}$, $\{n_\alpha\}$, $\{r_\alpha\}$, $\{s_\alpha\}$ are the sequences of indices occurring as in Proposition 6, then there classes ${}^*m$, ${}^*n$, ${}^*r$, ${}^*s$ are totally ordered in ${}^*\mathbb{Z}$, hence we may assume the representative sequences are. In particular, we may proceed with the same argument as in Proposition 6: the sequences $\{x_\alpha\}$, $\{x'_\alpha\}$ define elements of $[\![\widetilde{\mathscr{L}_\infty}]\!]$ which may be used to show that the composition $I_{{}^*v}\circ I_{{}^*w}$ is not defined. Now suppose that ${}^*v\notin{}^*V_\infty$ but ${}^*w\in{}^*V_\infty$. This means that both components of ${}^*v$ have the same order denoted $\mathsf{ord}_2({}^*v)$. Then there exists ${}^*x\in{}^*V_\infty$ such that ${}^*w+{}^*x\in{}^*V_\infty$ in which the two components of ${}^*w+{}^*x$ have order greater than $\mathsf{ord}_2({}^*v)$. Then both components of ${}^*v+{}^*w+{}^*x$ have equal order, which implies that $I_{{}^*v}\circ I_{{}^*w}$ is not defined. The case where ${}^*v\in{}^*V_\infty$ but ${}^*w\notin{}^*V_\infty$ is handled similarly. Now suppose ${}^*v,{}^*w\notin{}^*V_\infty$. Here there are two subcases. First suppose that the orders of the components of ${}^*v$, ${}^*w$ are not equal. Denote by $\mathsf{ord}_2({}^*v)$, $\mathsf{ord}_2({}^*w)$ the common order of the components of ${}^*v$, ${}^*w$. Then if say $\mathsf{ord}_2({}^*v)<\mathsf{ord}_2({}^*w)$, we define ${}^*x=(0,{}^*w_2)$ where ${}^*w_2$ is the second component of ${}^*w$. Then $I_{{}^*w}({}^*x)$ is defined but $I_{{}^*v+{}^*w}({}^*x)$ is not. If $\mathsf{ord}_2({}^*v)>\mathsf{ord}_2({}^*w)$ then $I_{{}^*v}$ is defined on ${}^*y=(0,{}^*v_2)$ but ${}^*y-{}^*w$ does not define an element of $\mathsf{Dom}(I_{{}^*w})$ since it does not converge to the same point in $[\![\widetilde{\mathscr{L}_\infty}]\!]$ as ${}^*y$. What remains is the case when $\mathsf{ord}_2({}^*v)=\mathsf{ord}_2({}^*w)$. If ${}^*v+{}^*w$ lies in ${}^*V_\infty$ then ${}^*w\in\mathsf{Dom}(I_{{}^*v})$ but not in $\mathsf{Ran}(I_{{}^*w})$. Otherwise, if the norms of the components of ${}^*v+{}^*w$ are equal, then $\mathsf{ord}_2({}^*v+{}^*w)>\mathsf{ord}_2({}^*v)=\mathsf{ord}_2({}^*w)$. If we let ${}^*x=(({}^*v+{}^*w)_1,0)$ then ${}^*x\in\mathsf{Dom}(I_{{}^*v+{}^*w})$ but not in $\mathsf{Dom}(I_{{}^*w})$ so it cannot be that $I_{{}^*v}\circ I_{{}^*w}=I_{{}^*v+{}^*w}$. □

Theorem 8. *As a set*

$$[\![\pi]\!]_1(\mathscr{L}_\infty,0,\mathsf{f}) \;=\; {}^*F_3\times[\![G]\!].$$

The composition ${}^\bullet u\circ{}^\bullet v$*, where* ${}^\bullet v=({}^*x,I_{{}^*v})$, ${}^\bullet w=({}^*y,I_{{}^*w})$ *is defined if and only if* ${}^*v=-{}^*w$.

Proof. Every element $\widetilde{\mathbb{I}}$ may be written in the form $I_v\circ\gamma=\gamma\circ I_v$ for $v\in V_\infty$ and $\gamma\in F_3$. Moreover, if $\widetilde{I}\in\widetilde{G}_n$, then $I\in G_n$. The second statement follows immediately from Proposition 7. □

Thus, although ${}^*F_3\times[\![G]\!]$ is formally a group, $[\![\pi]\!]_1(\mathscr{L}_\infty,0,\mathsf{f})$ is not a group with respect to the groupoid structure defined by its action on $[\![\widetilde{\mathscr{L}_\infty}]\!]$. It has nevertheless a distinguished subgroup isomorphic to ${}^*F_3$. On the other hand,

Theorem 9. *Any two elements* ${}^\bullet v$ *and* ${}^\bullet w$ *of* $[\![\pi]\!]_1(\mathscr{L}_\infty,0,\mathsf{f})$ *define composable elements of the mother germ* $[\![\pi]\!]_1(\mathscr{L}_\infty)$ *by restriction of domains.*

Proof. Let $I_{{}^*v}$, $I_{{}^*w}$ be the $[\![G]\!]$-coordinates of ${}^\bullet v$, ${}^\bullet w$. If ${}^*v$, ${}^*w$ and ${}^*v+{}^*w$ belong to ${}^*V_\infty$ then by restricting ${}^\bullet w$ to the leaf S_∞ and restricting ${}^\bullet v$ to the leaf of $[\![\widetilde{\mathscr{L}_\infty}]\!]$ containing ${}^\bullet w$, we obtain composable elements of $[\![\pi]\!]_1(\mathscr{L}_\infty)$. The other cases are handled similarly and are left to the reader. □

The lamination $\mathscr{L}_\infty$ has the following property: every leaf $L\neq S_\infty$ is conformal to either $\mathbb{C}$ or $\mathbb{C}^*=\mathbb{C}\setminus\{(0,0)\}$, [6]. Hence $\mathscr{L}_\infty$ is neither a suspension nor a locally free action of a Lie group. In particular, the antenna lamination is beyond the purview of the definition of $[\![\pi]\!]_1$ found in [4].

Given any leaf L of $\mathscr{L}_\infty$, one can obtain a graphical model T of L as a limit of a sequence of translations of T_∞. One can then repeat the discussion leading up to Theorem 8 for L. The proof of the following is left to the reader.

Theorem 10. *Let* $L\subset\mathscr{L}_\infty$ *be any leaf, modelled as above on a graph* $T\in\mathbb{A}$ *with vertex set* V. *Then for* $v\in V$ *and* f *constructed using a vector field as above,* $[\![\pi]\!]_1(\mathscr{L}_\infty,v,\mathsf{f})$ *may be identified with*

$${}^*\pi_1L\times[\![G]\!],$$

where ${}^*\pi_1 L \times \{0\}$ *is a subgroup with respect to the groupoid structure that is* $\cong 1$ *or* ${}^*\mathbb{Z}$.

Corollary 3. *Let* $v \in V_\infty \subset S_\infty$ *and* $v' \in V' \subset L' \neq S_\infty$. *Then choosing frame fields as above, the fundamental germs* $[\![\pi]\!]_1(\mathscr{L}_\infty, v, \mathsf{f})$ *and* $[\![\pi]\!]_1(\mathscr{L}_\infty, v', \mathsf{f}')$ *are not isomorphic.*

Proof. $[\![\pi]\!]_1(\mathscr{L}_\infty, v, \mathsf{f})$ has a nonabelian subgroup whereas $[\![\pi]\!]_1(\mathscr{L}_\infty, v', \mathsf{f}')$ is an abelian groupoid. □

8. THE $PSL(2,\mathbb{Z})$ ANOSOV FOLIATION

Let $\Gamma \subset PSL(2,\mathbb{R})$ be a discrete group of finite type, possibly with elliptic elements. The quotient $\Sigma = \mathbb{H}^2/\Gamma$ is a finite volume hyperbolic surface orbifold. The unit tangent bundle $T_1\Sigma$ is defined to be the quotient $\mathrm{T}_1\mathbb{H}^2/\Gamma$. Let $\rho : \Gamma \to \mathsf{Homeo}(\mathbb{S}^1)$ be the representation obtained by extending the action of Γ to the boundary of $\mathbb{H}^2$. Then $\mathrm{T}_1\Sigma$ may be identified with the suspension

$$\left(\mathbb{H}^2 \times \mathbb{S}^1\right)\Big/\Gamma$$

as follows. Given $(\tilde{z}, t) \in \mathbb{H}^2 \times \mathbb{S}^1$, associate $\mathsf{v}_{\tilde{z}} \in \mathrm{T}_1\mathbb{H}^2$, the vector based at $\tilde{z}$ and tangent to the ray limiting to t. This association is Γ-equivariant and descends to the desired homeomorphism. The expression of $\mathrm{T}_1\Sigma$ as a suspension defines a hyperbolic Riemann surface foliation $\mathscr{F}$ on $\mathrm{T}_1\Sigma$, which is also a fiber bundle over Σ provided that Γ has no elliptic points. $\mathscr{F}$ is called an *Anosov foliation*.

In [4], we worked with a definition of $[\![\pi]\!]_1$ that was available for suspensions such as $\mathscr{F}$ formed from fixed point free Γ. Unfortunately, this hypothesis excluded the most "explicit" of discrete subgroups of $PSL(2,\mathbb{R})$, the modular group $\Gamma = PSL(2,\mathbb{Z})$. The definition provided in this paper is clearly available in this case, and we devote the rest of this section to its consideration.

Two elements $r, s \in \mathbb{R} \cup \{\infty\} \approx \mathbb{S}^1$ are called *equivalent* if there exists $A \in PSL(2,\mathbb{Z})$ such that $A(r) = s$. Every equivalence class $[r]$ of extended reals corresponds to a leaf $L_{[r]}$ of $\mathscr{F}$, and since all $PSL(2,\mathbb{Z})$-orbits in $\mathbb{S}^1$ are dense, all leaves are dense. If $L_{[r]}$ is isomorphic to the punctured hyperbolic disk $\mathbb{D}^*$, then $[r]$ is quadratic over $\mathbb{Q}$. Otherwise, $L_{[r]}$ is isomorphic to $\mathbb{H}^2$.

Let us consider the leaf $L = L_{[0]} \cong \mathbb{D}^*$ covered by $\mathbb{H} \times \{0\}$. Choose $x \in L$ and a transversal T through x that is a fiber with respect to the projection onto the modular surface Σ. We assume that the lift $\tilde{x}$ of x to $\mathbb{H}^2$ is not an elliptic point for the action of $PSL(2,\mathbb{Z})$. Define f to be the lift of a frame on Σ based at the projection of T. As before, we denote by $\tilde{\mathsf{f}}$ the lift of f to $\widetilde{T}_0 \subset \mathbb{H}^2$ and by $\tilde{\mathsf{f}}_{\tilde{y}}$ its value at $\tilde{y} \in \widetilde{T}_0$. Note that for $A \in PSL(2,\mathbb{R})$, $A_*\tilde{\mathsf{f}}_{\tilde{x}} = A_*\tilde{\mathsf{f}}_{\tilde{y}}$ if and only if $A \in \Gamma = PSL(2,\mathbb{Z})$. A sequence $\{A_\alpha\} \subset PSL(2,\mathbb{Z})$ defines an f-diophantine approximation $\Leftrightarrow$ $(A_\alpha\tilde{x}, 0)$ projects to a sequence in T converging to x $\Leftrightarrow$ $(\tilde{x}, A_\alpha^{-1}(0))$ projects to a sequence in T converging to $x \Leftrightarrow A_\alpha^{-1}(0) \to 0$ in $\mathbb{S}^1$. Note that for any sequence $\{\gamma_{n_\alpha}\}$ in the deck group of $\mathbb{H}^2 \to L$,

$$\left\{ \gamma_n = \begin{pmatrix} 1 & 0 \\ n & 1 \end{pmatrix} \;\middle|\; n \in \mathbb{Z} \right\},$$

the sequence $\{\gamma_\alpha A_\alpha\}$ also defines an f-diophantine approximation. The fundamental germ $[\![\pi]\!]_1(\mathscr{F}, x, \mathsf{f})$ is then formed from the associated sequences $\{A_\alpha B_\alpha^{-1}\}$ where $\{B_\alpha\}$ is another f-diophantine approximation.

Note 3. The sequences $\{b_\alpha/d_\alpha = A_\alpha^{-1}(0)\}$ are *hyperbolic diophantine approximations*, as defined for example in [1]. See [4] for more on this point. In the case at hand, they give bad diophantine approximations of 0 whenever $b_\alpha \to \infty$, in the sense that it is never true that for some $c > 0$ and almost all α,

$$\left| 0 - \frac{b_\alpha}{d_\alpha} \right| \; < \; \frac{c}{d_\alpha^2}.$$

The f-diophantine approximations are not stable with respect to the operation of inversion. Indeed, let $r \in \mathbb{R}$ be any real number, $\{m_\alpha/n_\alpha\}$ a sequence of rationals (written in lowest terms) converging to r. Let M_α, N_α be such that $m_\alpha M_\alpha - n_\alpha N_\alpha = 1$. Assume that the indexing is such that $\alpha + N_\alpha/m_\alpha \to \infty$ as $\alpha \to \infty$. Then the sequence $\{X_\alpha\}$,

$$X_\alpha \; = \; \begin{pmatrix} -(\alpha m_\alpha + N_\alpha) & -m_\alpha \\ \alpha n_\alpha + M_\alpha & n_\alpha \end{pmatrix} \; \in \; PSL(2,\mathbb{Z}), \tag{5}$$

satisfies $X_\alpha^{-1}(0) \to 0$, but $X_\alpha(0) \to -r$. Using this fact, we can now show

Theorem 11. *As a set,*

$$[\![\pi]\!]_1(\mathscr{F}, x, \mathsf{f}) \; = \; PSL(2, {}^*\mathbb{Z}).$$

Proof. Let $\{A_\alpha\}$ be any sequence in $PSL(2,{}^*\mathbb{Z})$. Then after passing to a subsequence if necessary, we find $A_\alpha^{-1}(0) \to r$ for some $r \in \mathbb{R} \cup \{\infty\}$. Note that r is independent of the class of $\{A_\alpha\}$ in $PSL(2,{}^*\mathbb{Z})$. We may choose $\{n_\alpha\} \subset \mathbb{Z}$ so that $\gamma_{n_\alpha} A_\alpha^{-1}(0) \to 0$. Hence

$$\{A_\alpha\} \; = \; \{A_\alpha \gamma_{-n_\alpha}\} \cdot \{\gamma_{-n_\alpha}^{-1}\}$$

defines an element of $[\![\pi]\!]_1(\mathscr{F}, x, \mathsf{f})$. $\square$

It is not difficult to see that with respect to its action on the germ universal cover $[\![\widetilde{\mathscr{F}}]\!]$, $[\![\pi]\!]_1(\mathscr{F}, x, \mathsf{f})$ is not a group. Indeed, the class of the sequence $\{X_\alpha^{-1}\}$, where $\{X_\alpha\}$ is the sequence appearing in (5), is not defined on $\widetilde{L}$.

9. MAPPING CLASS GROUP OF THE ALGEBRAIC UNIVERSAL COVER OF A SURFACE

In this section, we use the fundamental germ to prove a Nielsen type theorem for the algebraic universal cover of a closed surface. We begin by recalling a few facts, referring the reader to [5] for details.

Let Σ be a closed surface and let $\mathscr{G} = \{G_\alpha\}$ be the set of all normal finite index subgroups. For each G_α, there exists a covering $\sigma_\alpha : \Sigma_\alpha \to \Sigma$ defined by the condition that $\pi_1\Sigma_\alpha$ maps isomorphically onto G_α. If $G_\alpha \subset G_\beta$, there is a unique covering $s_{\alpha\beta} : \Sigma_\alpha \to \Sigma_\beta$ for which $\sigma_\alpha = \sigma_\beta \circ s_{\alpha\beta}$. Hence the collection of σ_α and $s_{\alpha\beta}$ forms an inverse system of surfaces by covering maps.

Definition 6. The **algebraic universal cover** of Σ is the inverse limit

$$\widehat{\Sigma} \; = \; \varprojlim \Sigma_\alpha.$$

If $\sigma : Z \to \Sigma$ is any finite covering, then σ lifts to a homeomorphism

$$\hat{\sigma} : \widehat{Z} \to \widehat{\Sigma}.$$

Thus the algebraic universal cover depends only on the type of Σ (elliptic, parabolic, hyperbolic). In fact, there are only two non-trivial examples of algebraic universal covers of closed surfaces: that of the torus and that of a surface of hyperbolic type.

The inverse limit

$$\hat{\pi}_1\Sigma \; = \; \varprojlim \big(\pi_1\Sigma\big)/G_\alpha$$

is a Cantor group called the *profinite completion* of $\pi_1\Sigma$. The homomorphism $i : \pi_1\Sigma \rightarrow \hat{\pi}_1\Sigma$ induced by the system of projections $\pi_1\Sigma \longrightarrow \pi_1\Sigma/G_\alpha$ has dense image. Define a representation

$$\varsigma : \pi_1\Sigma \longrightarrow \mathsf{Homeo}(\hat{\pi}_1\Sigma)$$
$$\varsigma_\gamma(\hat{g}) = \hat{g}\cdot i(\gamma)^{-1}$$

for $\gamma \in \pi_1\Sigma$ and $\hat{g} \in \hat{\pi}_1\Sigma$. Then we may identify $\widehat{\Sigma}$ with the suspension of ς:

$$\widehat{\Sigma} \;\approx\; \big(\widetilde{\Sigma}\times\hat{\pi}_1\Sigma\big)\big/\pi_1\Sigma.$$

With this identification, we see that $\widehat{\Sigma}$ is a surface lamination with Cantor transversals homeomorphic to $\hat{\pi}_1\Sigma$, that is, a solenoid. Moreover, it can also be seen from this presentation that every leaf L of $\widehat{\Sigma}$ satisfies

$$\pi_1 L \;\cong\; \bigcap G_\alpha.$$

However for closed surfaces, $\bigcap G_\alpha = 1$, so here, L is simply connected. Each leaf L is dense and a path-component of $\widehat{\Sigma}$. For every α, the pre-image of the projection map $\widehat{\Sigma} \rightarrow \Sigma_\alpha$ is a fiber transversal, homeomorphic to $\hat{\pi}_1\Sigma_\alpha \cong \widehat{G}_\alpha$.

Now let L be a fixed leaf of $\widehat{\Sigma}$.

Definition 7. The **leafed mapping class group** of $\widehat{\Sigma}$ is

$$\mathsf{MCG}(\widehat{\Sigma},L) \;=\; \mathsf{Homeo}(\widehat{\Sigma},L)/\simeq,$$

where $\simeq$ is the relation of homotopy of homeomorphisms.

We denote by $[h]$ the mapping class associated to a homeomorphism h.

Let G be a group.

Definition 8. The **virtual automorphism group** of G is

$$\mathsf{Vaut}(G) \;=\; \Big\{\phi : H \rightarrow H' \;\Big|\; \phi \text{ an isomorphism and } H, H' \text{ finite index subgroups of } G\Big\}\Big/\sim,$$

where $\phi_1 \sim \phi_2$ if there exists $H'' < G$ of finite index, contained in $\mathsf{Dom}(\phi_1)\cap\mathsf{Dom}(\phi_2)$ and such that $\phi_1|_{H''} = \phi_2|_{H''}$.

Note that the equivalence relation $\sim$ is precisely what is needed to make composition of virtual automorphisms well-defined. We point out also that if $H < G$ is of finite index, then $\mathsf{Vaut}(H) \cong \mathsf{Vaut}(G)$.

Theorem 12. $\mathsf{MCG}(\widehat{\Sigma},L) \;\cong\; \mathsf{Vaut}(\pi_1\Sigma)$.

Note 4. We first learned the statement of this theorem in a conversation with D. Sullivan in 1995. The first proof appeared in the thesis of C. Odden [8].

Proof. Define a homomorphism

$$\Theta : \mathsf{Vaut}(\pi_1\Sigma) \longrightarrow \mathsf{MCG}(\widehat{\Sigma},L)$$

as follows. Given $\phi : G_1 \rightarrow G_2$ an isomorphism of finite index subgroups of $\pi_1\Sigma$, we may find covers $\sigma_1, \sigma_2 : \Sigma' \rightarrow \Sigma$ – indexed by G_1 and G_2 – so that

$$(\sigma_2)_* \circ (\sigma_1)_*^{-1} \;=\; \phi\,:$$

this follows from the classical Nielsen theorem. Then we define

$$\Theta(\phi) \;=\; [\hat{\sigma}_2 \circ \hat{\sigma}_1^{-1}]$$

where for $i = 1,2$, $\hat{\sigma}_i : \widehat{\Sigma}' \to \widehat{\Sigma}$ is the algebraic universal cover lift. If $G' < \mathsf{Dom}(\phi)$, then $\Theta(\phi|_{G'}) \;=\; \Theta(\phi)$, since $\Theta(\phi|_{G'})$ is defined by the pair $\sigma_i \circ \sigma$, $i = 1,2$, where $\sigma : \Sigma'' \to \Sigma'$ is a cover for which $\sigma_1 \circ \sigma$ is indexed by G'. Thus Θ is a well-defined homomorphism.

Claim 1. *Θ is onto.*

Let $h : (\widehat{\Sigma}, L) \to (\widehat{\Sigma}, L)$ be a homeomorphism. After performing an isotopy, we may arrange that h fixes a point x and fiber transversal T containing x. Without loss of generality, we may assume that T is a fiber transversal over Σ. Due to the suspension structure, $T \approx \hat{\pi}\Sigma$: fix this identification so that $x \mapsto 1$ and $L \cap T \mapsto \pi_1\Sigma$. Since $h(L) = L$, we obtain a bijection

$$h_* : \pi_1\Sigma \longrightarrow \pi_1\Sigma$$

in which $h(1) = 1$.

Suppose that for each $G_\alpha < \pi_1\Sigma$, $h_*|G_\alpha$ is not homomorphic. This means that for every α, there exists $\gamma_\alpha, \gamma'_\alpha \in G_\alpha$ so that

$$h_*\big(\gamma_\alpha \cdot \gamma'_\alpha\big) \neq h_*\big(\gamma_\alpha\big) \cdot h_*\big(\gamma'_\alpha\big). \tag{6}$$

Assuming that $\widehat{\Sigma}$ has been equipped with a hyperbolic metric, say lifted from Σ, then the sequences $\{\gamma_\alpha\}$, $\{\gamma'_\alpha\}$ define elements of the fundamental germ

$$^*\gamma,\ ^*\gamma' \;\in\; [\![\pi]\!]_1(\widehat{\Sigma}, x, \mathsf{f})$$

where f is a frame field lifted from a frame on Σ. But this fundamental germ is a subgroup of the mother germ $[\![\pi]\!]_1(\widehat{\Sigma})$. By Theorem 7, h induces a groupoid isomorphism

$$[\![h]\!]_* : [\![\pi]\!]_1(\widehat{\Sigma}) \;\longrightarrow\; [\![\pi]\!]_1(\widehat{\Sigma}),$$

and so we must have

$$[\![h]\!]_*\big(^*\gamma \cdot {}^*\gamma'\big) \;=\; [\![h]\!]_*\big(^*\gamma\big) \cdot [\![h]\!]_*\big(^*\gamma'\big).$$

This contradicts equation (6). Thus h_* defines an isomorphism when restricted to some G_α, and this isomorphism determines an element $\phi \in \mathsf{Vaut}(\pi_1\Sigma)$. Note that that ϕ does not depend on the isotopy used to ensure $h(x) = x$ since the holonomy group of $\widehat{\Sigma}$ at any point is trivial. Choose $\sigma_i : \Sigma' \to \Sigma$, $i = 1,2$, so that $\Theta(\phi) = [\hat{\sigma}_2 \circ \hat{\sigma}_1^{-1}]$. To simplify notation, we write $h_0 = \hat{\sigma}_2 \circ \hat{\sigma}_1^{-1}$.

Recall that since $\widehat{\Sigma}$ is compact with hyperbolic leaves, the germ universal cover of $\widehat{\Sigma}$ is $^\bullet\mathbb{H}^2$. The homeomorphisms h and h_0 lift to the standard bijections $^\bullet h$ and $^\bullet h_0$ of $^\bullet\mathbb{H}^2$ sharing the same equivariance with respect to the action of the mother germ $[\![\pi]\!]_1(\widehat{\Sigma})$. In particular, they act identically on the set of galaxies of $^\bullet\mathbb{H}^2$. For this reason, we may choose a germ universal cover topology for $^\bullet\mathbb{H}^2$ with respect to which both $^\bullet h$ and $^\bullet h_0$ are homeomorphisms.

Define a homotopy $^\bullet H_t$ from $^\bullet h$ to $^\bullet h_0$ as follows. For each $^\bullet z \in {}^\bullet\mathbb{H}^2$, $^\bullet H_t\big(^\bullet z\big)$ is the point subdividing the hyperbolic geodesic connecting $^\bullet h\big(^\bullet z\big)$ to $^\bullet h_0\big(^\bullet z\big)$ into the proportion $t : 1 - t$. By construction, $^\bullet H_t$ has the same equivariance as $^\bullet h$ and $^\bullet h_0$ and is in particular continuous. Since its initial and final maps are standard, so is $^\bullet H_t$. By Proposition 6, it descends to a homotopy H_t of h and h_0. It follows that $[h] = [h_0] = \Theta(\phi)$, and Θ is onto.

Claim 2. Θ is one-to-one.

If not, then there exists $\phi \neq$ the identity map with $\Theta(\phi) = 1$. But then $\Theta(\phi)$ would have to induce the identity map on the mother germ; by construction, this can only happen if ϕ is trivial. □

Let $\mathsf{Mod}(\widehat{\Sigma},L)$ be the *Teichmüller modular group* of the pair $(\widehat{\Sigma},L)$: the group of homotopy classes of quasiconformal homeomorphisms of $\widehat{\Sigma}$ that preserve L.

Corollary 4. $\mathsf{Mod}(\widehat{\Sigma},L)=\mathsf{MCG}(\widehat{\Sigma},L)$.

Proof. This follows from the proof of Theorem 12 and the fact that every finite cover of compact Riemann surfaces is homotopic to a quasiconformal cover. □

Theorem 12 can be used to formulate the following conjectural Nielsen-type theorem. Given $x\in L$, the fundamental germ $[\![\pi]\!]_1(\widehat{\Sigma},x)$ is made up of all sequences $\{\gamma_\alpha\}$ converging with respect to the lattice of finite index normal subgroups G of $\pi_1\Sigma$, so

$$[\![\pi]\!]_1(\widehat{\Sigma},x) \;\cong\; \bigcap_{[\pi_1\Sigma:G]<\infty} {}^*G \;\subset\; {}^*\pi_1\Sigma.$$

It follows then that there is a monomorphism $\mathsf{Vaut}(\pi_1)\hookrightarrow\mathsf{Aut}([\![\pi]\!]_1(\widehat{\Sigma},x))$, which descends to $\mathsf{Vaut}(\pi_1)\hookrightarrow\mathsf{Out}([\![\pi]\!]_1(\widehat{\Sigma},x))$ upon passage to the quotient. This latter map is also a monomorphism, since no nontrivial virtual automorphism ϕ can induce on $[\![\pi]\!]_1(\widehat{\Sigma},x)$ an inner automorphism. For otherwise, ϕ would have to be inner on some subgroup H, hence all, which is only possible if ϕ is trivial. In view of these remarks we

Conjecture. *The monomorphism* $\mathsf{MCG}(\widehat{\Sigma},L)\hookrightarrow\mathsf{Out}([\![\pi]\!]_1(\widehat{\Sigma},x))$ *is an isomorphism.*

We end this section by explaining the importance of Theorem 12 and Corollary 4 in giving a genus independent reformulation the Ehrenpreis conjecture. The classical Ehrenpreis conjecture is:

> *Given two closed hyperbolic surfaces* Σ_1 *and* Σ_2 *and* $\varepsilon>0$, *there exist finite, locally isometric covering surfaces* Z_1 *and* Z_2 *of each which are* $(1+\varepsilon)$*-quasiisometric.*

We then have the following equivalent, genus independent version:

> *Every orbit of the action of* $\mathsf{Mod}(\widehat{\Sigma},L)$ *on* $\mathscr{T}(\widehat{\Sigma})$ *is dense.*

In other words, the genus independent version says that, although the moduli space

$$\mathscr{T}(\widehat{\Sigma})/\mathsf{Mod}(\widehat{\Sigma},L)$$

is uncountable, it has the "topology of a point" (*i.e.* the coarse topology). If affirmed, the Ehrenpreis conjecture would thus provide an explanation for the jump between the existence of moduli (dimension 2) and rigidity (dimension 3 and higher) in hyperbolic geometry. See the articles [2], [5] for more discussion.

REFERENCES

[1] Bernik, V. I. & Dodson, M. M., *Metric Diophantine Approximation on Manifolds.* Cambridge Tracts in Mathematics **137**. Cambridge University Press, Cambridge, 1999.

[2] Biswas, I., Nag, S. & Sullivan, D.P., Determinant bundles, Quillen metrics and Mumford isomorphisms over the universal commensurability Teichmüller space. *Acta Math.* **176** (1996), no. 2, 145–169.

[3] Gendron, T. M., *Fuchsian Germs.* PhD Thesis, City University of New York, 1997.

[4] Gendron, T. M., *The algebraic theory of the fundamental germ.* To appear in the Bulletin of the Brazilian Mathematical Society, math.DG/0506270.

[5] Gendron, T. M., *The Ehrenpreis conjecture and the moduli-rigidity gap.* in "Complex Manifolds and Hyperbolic Geometry" (eds. Earle et al), 207–229, *Contemporary Mathematics*, AMS, Providence, RI, 2002.

[6] Ghys, E., *Laminations par surfaces de Riemann.* in "Dynamique et géométrie complexes (Lyon, 1997)", 49–95, *Panoramas & Synthèse* **8**, Soc. Math. France, Paris, 1999.

[7] Goldbladt, R. *Lectures on the Hyperreals.* Springer-Verlag, New York, NY, 1998.

[8] Odden, C., *PhD Thesis*, Duke University, 1997.

[9] Robinson, A., *Non-standard Analysis*, Princeton Landmarks in Mathematics. Princeton University Press, Princeton, NJ, 1996.

[10] Stroyan, K. D., Luxemburg, W. A. J. *Introduction to the theory of infinitesimals.* Pure and Applied Mathematics, No. 72. Academic Press, New York-London, 1976.

INSTITUTO DE MATEMÁTICAS – UNIDAD CUERNAVACA, UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO, AV. UNIVERSIDAD S/N, C.P. 62210 CUERNAVACA, MORELOS, MÉXICO

E-mail address: `tim@matcuer.unam.mx`